

#### RayBiotech, Inc.

3607 Parkway Lane suite 200 Norcross,GA 30092 Tel: 770-729-2992, 1-888-494-8555 Fax: 770-206-2393

Website: www.raybiotech.com Email: info@raybiotech.com

## Certificate of Analysis and Data Sheet

# Rabbit anti-Human Cardiotrophin-1 antibody

Catalog No:	= Isotype:	= Speceis:	Accession No:	
130-10013	Rabbit Ig G	Human	Q16619	

#### Description


Cardiotrophin-1 (CT-1) is a cardiac hypertrophic factor and belongs to the IL-6 cytokine family. The protein interacts with the glycoprotein 130 (gp130)/leukemia inhibitory factor receptor beta (LIFR) heterodimer. CT-1 deficiency causes increased motoneuron cell death in spinal cord and brainstem nuclei of mice. In addition, CT-1 enhances transcription factor NF kappa B DNA-binding activities and activates phosphatidylinositol 3-kinase (PI-3 kinase) in cardiac myocytes. Studies have revealed CT-1 is highly expressed in the heart, prostate, skeletal muscle, and ovary, and expressed at the lower levels in lung, kidney, pancreas, thymus, small intestine, and testis. CT-1 is highly secreted by myotubes, and promotes the survival of cultured embryonic mouse and rat MNs. CT-1 is associated with pathophysiology of heart diseases, like hypertension, myocardial infarction, valvular heart disease, and congestive heart failure.

### **Applications**

Table Summary of antibody applications and working conditions

Options Functions	YES	NO	Not determined	Recommended Work dilution or concentration
ELISA	*			1:80000(at least detecting 12.5ng/ml)
Western Blotting	*			1:1000-5000(at least)
Enzyme Immunoassay(EIA)			#	

Note: Other applications are not tested yet. Optimal dilutions should be determined by each laboratory for each application



Immunodetection Analysis: Representative blot from a previous lot. Lane 1.recombinant protein CT-1. Lane 2. HepG2 cell lysate. The membrane blot was probed with anti-CT-1 primary antibody(0.25μg/ml). Proteins were visualized using a goat anti-rabbit secondary antibody conjugated to HRP and chemiluminescence detection system. Arrows indicate cellular CT-1 from human and mouse cells (24 kDa).

The products are furnished for LABORATORY RESEARCH USE ONLY.

Not for diagnostic or therapeutic use.



#### RayBiotech, Inc.

3607 Parkway Lane suite 200 Norcross,GA 30092 Tel: 770-729-2992, 1-888-494-8555 Fax: 770-206-2393

Website: www.raybiotech.com Email: info@raybiotech.com

#### Preparation

Immunogen was expressing Human CT-1 protein. This antibody was produced from a rabbit immunized with the immunogen. The IgG fraction was purified from rabbit serum followed by Protein A/G affinity chromatography.

### Specificity

Rabbit anti-Human Cardiotrophin-1 antibody detects specifically target derived from human recombinant protein as well as HepG2 cell lysates. Cross reactivity with mouse and rat CT-1 was not tested yet.

#### Reconstitution

Supplied as lyophilized and purified antibody originally containing PBS, without **Preservative Stabilizers**, liking Sodium Azide. *It final concentration is indicated in shipping vial*. **Please avoid freeze-thaw cycles**.

The antibody is stable for at least years from the data of receipt when stored at  $-20^{\circ}$ C to  $-70^{\circ}$ C. Reconstituted antibody (suggesting with sterile PBS) can also be aliquotted and stored frozen at  $-20^{\circ}$ C to  $-70^{\circ}$ C in a manual defrost freezer for months without detectable loss activity. Upon reconstitution, the antibody can be stored at  $4^{\circ}$ C for 1 month.

### Storage

Store at -20°C, if not intended for use within a month, keep it at 4°C and minimize freezing and thawing when use.

## Related products

- 1. Human Cardiotrophin-1 (CT-1) ELISA Kit For Serum, Plasma, Cell Culture Supernatant and Urine (Cat# ELH-CT1-001)
- 2. Recombinant Human Cardiotrophin-1 (Cat# 230-00008)

#### Reference

- 1. Arce V, et. al. (1998) Synergistic effects of Schwann- and muscle-derived factors on motoneuron survival involve GDNF and cardiotrophin-1 (CT-1). *J Neurosci* 18:1440-1448.
- 2. Ronald W, et al. (2001). Cardiotrophin-1, a muscle-derived Cytokine, is required for the survival of subpopulations of developing motoneurons. *J Neurosci.* 21:1283-1291.
- 3. Arce V, et al. (1999) Cardiotrophin-1 requires LIFRbeta to promote survival of mouse motoneurons purified by a novel technique. *J Neurosci Res* 55:119-126.

The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.