



RayBiotech, Inc.

3607 Parkway Lane suite 200

Norcross, GA 30092

Tel: 770-729-2992, 1-888-494-8555

Fax: 770-206-2393

Website: www.raybiotech.com

Email: info@raybiotech.com

Recombinant Human S100 Calcium Binding Protein B (S100B)

Catalog No.	Size	Species	Protein Accession No.
230-00002	10, 50, 100 µg	Human	P04271

Synonyms

S100; S100beta; S-100 calcium-binding protein, beta chain; S100 calcium binding protein B.

Description

S100B protein plays an important role in neurite extension, proliferation of melanoma cells, stimulation of Ca^{2+} fluxes, inhibition of PKC-mediated phosphorylation, astrocytosis and axonal proliferation, and inhibition of microtubule assembly. S100B is glial-specific and is expressed primarily by a subtype of mature astrocytes that ensheath blood vessels and by NG2-expressing cells. Chromosomal rearrangements and altered expression of this gene have been implicated in several neurological, neoplastic, and other types of diseases, including Alzheimer's disease, Down's syndrome, epilepsy, amyotrophic lateral sclerosis, melanoma, and type I diabetes. It is found that S100B interacts with S100A11, IQGAP1, S100 calcium binding protein A1, Tau protein, S100A6, P53, etc.

Preparation

The full-length of human S100B gene was cloned and expressed in *Escherichia coli*. The recombinant protein has an N-terminal 6×histidine tag and was purified by immobilized metal ion affinity chromatography (IMAC).

Source

Recombinant histidine-tagged protein, purified from *E. coli*.

Predicted Molecular Mass

~12.5 kDa with the 6×histidine tag.

Formulation

Fine white powder, lyophilized. S100B protein was constituted in a 0.2 µm filtered phosphate-buffered saline (1x PBS) before lyophilized.

Solubility

The product is soluble in the neutral buffer indicated above.

Storage

The protein is stable at -20 to -70 °C in a manual defrost freezer. **Avoid repeated freeze-thaw cycles.**

Purity

>95%, determined by SDS-PAGE and stained with Coomassie blue.

Applications

	Yes	No	Not Determined
ELISA	√		
Western blotting	√		
Dot blotting	√		
Protein array	√		
Others			√

Note: other applications are not tested yet. Optimal working conditions should be determined by each laboratory for each application.

References

1. Mbele, GO, et al. (2002). The zinc- and calcium-binding S100B interacts and co-localizes with IQGAP1 during dynamic rearrangement of cell membranes. *J. Biol. Chem.* **277** (51): 49998–50007.
2. Fackler, OT, et al. (1999). Activation of Vav by Nef induces cytoskeletal rearrangements and downstream effector functions. *Mol. Cell.* **3** (6): 729–39.
3. Gentil, BJ, et al. (2001). The giant protein AHNK is a specific target for the calcium- and zinc-binding S100B protein: potential implications for Ca^{2+} homeostasis regulation by S100B. *J. Biol. Chem.* **276** (26): 23253–61.
4. Lin, Jing, et al. (2004). Inhibiting S100B restores p53 levels in primary malignant melanoma cancer cells. *J. Biol. Chem.* **279** (32): 34071–7.

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**