

Recombinant Human Sonic Hedgehog (H-SHH)

CATALOG #: 4010-25 25 μg

4010-100 100 μg 4010-1000 1 mg

LOT #: _____

SOURCE: E. coli

PURITY: > 97% by SDS-PAGE and HPLC analyses

Endotoxin level is < 0.01 ng per µg rh-SHH

MOLECULAR WT: 19.697 kDa

FORM: Lyophilized from a 10 mM sodium phosphate buffer, pH 7.5.

RECONSTITUTION:

Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water to a concentration of 0.1-0.5 mg/ml, which can be further diluted into other aqueous solutions.

STORAGE CONDITIONS:

The lyophilized human Sonic Hedgehog is very stable at -20 °C. Upon reconstitution the material should be aliquoted and frozen at -20 °C. It is recommended to add a carrier protein (0.1% HSA or BSA) for long term storage.

DESCRIPTION:

Recombinant Human Sonic Hedgehog (SHH) is a highly conserved protein that plays an important role in embryonic development. It is expressed in neural tissue, the gut, and areas of limb development and promotes differentiation and growth in a tissue-specific manner. SHH is synthesized as a 45-kDa precursor protein, which is then cleaved to generate the active 19-kDa N-terminus. SHH interacts with the Patched and Smoothened transmembrane receptors, leading to the activation of GLI family transcription factors. Disruption of any part of this pathway during embryogenesis is associated with birth defects ranging from mild to severe. In adults, abnormal activation of the SHH pathway has been implicated in several forms of cancer. Three mammalian hedgehog genes (sonic, desert, Indian) share about 60% homology. The Cys at position 25 has been substituted with Ile. Human SHH is a non-glycosylated polypeptide consisting of 175 amino acids with a molecular weight of 19,697 Da.

BIOLOGICAL ACTIVITY:

Human recombinant Sonic Hedgehog is biologically active as compared to a standard. The ED₅₀, measured by the dose-dependent induction of alkaline phosphatase production by C3H/10T1/2 fibroblasts, is 750ng/ml.

AMINO ACID SEQUENCE:

IIGPGRGFG KRRHPKKLTP LAYKQFIPNV AEKTLGASGR YEGKISRNSE RFKELTPNYN PDIIFKDEEN TGADRLMTQR CKDKLNALAI SVMNQWPGVK LRVTEGWDED GHHSEESLHY EGRALDITTS DRDRSKYGML ARLAVEAGFD WVYYESKAHI HCSVKAENSV AAKSGG

FOR RESEARCH USE ONLY! Not to be used on humans.

RELATED PRODUCTS:

rev.02/11

Apoptosis Detection Kits & Reagents

- Annexin V Kits & Bulk Reagents
- Caspase Assay Kits & Reagents
- Mitochondrial Apoptosis Kits & Reagents
- Nuclear Apoptosis Kits & Reagents
- Apoptosis Inducers and Set
- Apoptosis siRNA Vectors

Cell Fractionation System

- Mitochondria/Cytosol Fractionation Kit
- Nuclear/Cytosol Fractionation Kit
- Membrane Protein Extraction Kit
- Cytosol/Particulate Rapid Separation Kit
- Mammalian Cell Extraction Kit
- FractionPREP Fractionation System

Cell Proliferation & Senescence

- Quick Cell Proliferation Assay Kit
- Senescence Detection Kit
- High Throughput Apoptosis/Cell Viability Assay Kits
- LDH-Cytotoxicity Assay Kit
- Bioluminescence Cytotoxicity Assay Kit
- Live/Dead Cell Staining Kit

Cell Damage & Repair

- HDAC Fluorometric & Colorimetric Assays & Drug Discovery Kits
- HAT Colorimetric Assay Kit & Reagents
- DNA Damage Quantification Kit
- Glutathione & Nitric Oxide Fluorometric & Colorimetric Assay Kits

Signal Transduction

- cAMP & cGMP Assay Kits
- Akt & JNK Activity Assay Kits
- Beta-Secretase Activity Assay Kit

Adipocyte & Lipid Transfer

- Recombinant Adiponectin, Survivin, & Leptin
- CETP Activity Assay & Drug Discovery Kits
- PLTP Activity Assay & Drug Discovery Kits
- Total Cholesterol Quantification Kit

Molecular Biology & Reporter Assays

- siRNA Vectors
- Cloning Insert Quick Screening Kit
- Mitochondrial & Genomic DNA Isolation Kits
- 5 Minutes DNA Ligation Kit
- 20 Minutes Gel Staining/Destaining Kit
- β -Galactosidase Staining Kit & Luciferase Reporter Assay Kit

Growth Factors and Cytokines

Monoclonal and Polyclonal Antibodies