Recombinant Human BD-4

CATALOG #: 4525-20 20 μg

4525-1000 1 mg

LOT #: _____

SYNONYMS: HBD-4, DEFB-4, Beta-defensin 4

SOURCE: E. coli

PURITY: >98% by SDS-PAGE and RP-HPLC analyses

FORM: Sterile filtered and lyophilized from 20 mM PBS, pH 7.4 and

130 mM NaCl.

RECONSTITUTION:

Centrifuge the vial prior to opening. Reconstitute in sterile ddH_2O to a concentration $\geq 100 \, \mu g/ml$. This solution can then be diluted into other aqueous buffers and stored at 4 $^{\circ}C$ for 2-7 days and at -20 $^{\circ}C$ for future use.

STORAGE CONDITIONS:

The lyophilized protein is best-stored desiccated at -20 °C. Reconstituted human BD-4 should be stored in working aliquots at -20 °C.

DESCRIPTION:

Defensins are cationic peptides with a large spectrum of antimicrobial activity that comprise an important arm of the innate immune system. The Alpha defensins are differentiated from the Beta-defensins by the pairing of their 3 disulfide bonds. Four human Beta-defensins have been identified to date; BD-1, BD-2, BD-3 and BD-4. Beta-defensins are expressed on some leukocytes and at epithelial surfaces. In addition to their direct antimicrobial activities, they are chemoattractant towards immature dendritic cells and memory T cells. Beta-defensin proteins are expressed as the C-terminal portion of precursors and are released by proteolytic cleavage of a signal sequence (for BD-1 a pro-peptide region). Beta-defensins contain a six-cysteine motif that forms three intra-molecular disulfide bonds. Beta Defensin-4 Human Recombinant produced in *E. coli* is a single, non-glycosylated polypeptide chain containing 50 amino acids. The BD-4 is purified by proprietary chromatographic techniques.

AMINO ACID SEQUENCE: (MW 6 kDa)

EFELDRICGY GTARCRKKCR SQEYRIGRCP NTYACCLRKW DESLLNRTKP

BIOLOGICAL ACTIVITY:

Determined by its ability to chemoattract human monocytes using a concentration range of 0.1-50 ng/ml.

RELATED PRODUCTS:

rev. 02/11

Apoptosis Detection Kits & Reagents

- Annexin V Kits & Bulk Reagents
- Caspase Assay Kits & Reagents
- Mitochondrial Apoptosis Kits & Reagents
- Nuclear Apoptosis Kits & Reagents
- Apoptosis Inducers and Set
- Apoptosis siRNA Vectors

Cell Fractionation System

- Mitochondria/Cytosol Fractionation Kit
- Nuclear/Cytosol Fractionation Kit
- Membrane Protein Extraction Kit
- Cytosol/Particulate Rapid Separation Kit
- Mammalian Cell Extraction Kit
- FractionPREP Fractionation System

Cell Proliferation & Senescence

- Quick Cell Proliferation Assay Kit
- Senescence Detection Kit
- High Throughput Apoptosis/Cell Viability Assay Kits
- LDH-Cytotoxicity Assay Kit
- Bioluminescence Cytotoxicity Assay Kit
- Live/Dead Cell Staining Kit

Cell Damage & Repair

- HDAC Fluorometric & Colorimetric Assays & Drug Discovery Kits
- HAT Colorimetric Assay Kit & Reagents
- DNA Damage Quantification Kit
- Glutathione & Nitric Oxide Fluorometric & Colorimetric Assav Kits

Signal Transduction

- cAMP & cGMP Assay Kits
- Akt & JNK Activity Assay Kits
- Beta-Secretase Activity Assay Kit

Adipocyte & Lipid Transfer

- Recombinant Adiponectin, Survivin, & Leptin
- CETP Activity Assay & Drug Discovery Kits
- PLTP Activity Assay & Drug Discovery Kits
- Total Cholesterol Quantification Kit

Molecular Biology & Reporter Assays

- siRNA Vectors
- Cloning Insert Quick Screening Kit
- Mitochondrial & Genomic DNA Isolation Kits
- 5 Minutes DNA Ligation Kit
- 20 Minutes Gel Staining/Destaining Kit
- β-Galactosidase Staining Kit & Luciferase Reporter Assay Kit

Growth Factors and Cytokines

Monoclonal and Polyclonal Antibodies

FOR RESEARCH USE ONLY! Not to be used on humans.

Tel: 408-493-1800 | Fax: 408-493-1801 www.biovision.com | tech@biovision.com