

### RayBiotech, Inc.

3607 Parkway Lane suite 200 Norcross,GA 30092 Tel: 770-729-2992, 1-888-494-8555

Fax: 770-206-2393

Website: www.raybiotech.com Email: info@raybiotech.com

## Certificate of Analysis and Data Sheet

# Rat Anti-Mouse CD200R With HRP-conjugated Secondary Antibody

Catalog No.Target SpeciesIsotypeDS-MB-00481MouseIgG2a

## **Preparation**

Synonyms: CD200R, CRTR2, MOX2R, OX2R

Purification: Purified IgG prepared by affinity chromatography on Protein G from tissue culture

supernatant

Immunogen: Fusion protein mCD200RCD4

## Specificity

DS-MB-00481 recognizes mouse CD200R, a cell surface glycoprotein (also known as OX2R), In mouse CD200R is expressed primarily by peripheral blood monocytes and neutrophils but also by other leucocytes including T-lymphocytes and mast cells. Studies suggest that CD200-CD200R interaction may be involved in the control of myeloid cellular function.

#### **Formulation**

**Product Type:** Monoclonal Antibody **Product Form:** Purified IgG - liquid **Buffer Solution:** Phosphate buffered saline

Preservative Stabilizers: 0.09% Sodium Azide (NaN<sub>3</sub>)

**Approx. Protein Concentrations:** IgG concentration 1.0 mg/ml

**Applications** 

Options Functions	YES	NO	Not determined	Recommended Work dilution or concentration
Flow Cytometry	•			1/25 - 1/50
ELISA			•	

Note: Other applications are not tested yet. Optimal dilutions should be determined.



### RayBiotech, Inc.

3607 Parkway Lane suite 200 Norcross, GA 30092 Tel: 770-729-2992, 1-888-494-8555

Fax: 770-206-2393

Website: www.raybiotech.com Email: info@raybiotech.com

## Secondary Antibody Applications

Options Functions	YES	NO	Not determined	Recommended Work dilution or concentration
Immunoassay (ELISA, Western blot)	•			1:5,000-1:10,000

## Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

**Shelf Life:** 18 months from date of dispatch.

#### References

- 1. Zhao, J. et al. (2009) Evasion by stealth: inefficient immune activation underlies poor T cell response and severe disease in SARS-CoV-infected mice. PLoS Pathog. 5: e1000636.
- 2. Björck, P. et al. (2011) Plasmacytoid Dendritic Cell Dichotomy: Identification of IFN-{alpha} Producing Cells as a Phenotypically and Functionally Distinct Subset. J Immunol. 186: 1477-85.
- 3. Dellacasa-Lindberg, I. et al. (2011) Migratory Activation of Primary Cortical Microglia upon Infection with Toxoplasma gondii. Infect Immun. 79: 3046-52.
- 4. Krejsek, J. et al. (2010) Expression of CD200/CD200R regulatory molecules on granulocytes and monocytes is modulated by cardiac surgical operation. Perfusion. 25: 389-97.
- 5. Rijkers, E.S. et al. (2008) The inhibitory CD200R is differentially expressed on human and mouse T and B lymphocytes. Mol Immunol. 45: 1126-35.
- 6. Costello, D.A. et al. (2011) Long Term Potentiation Is Impaired in Membrane Glycoprotein CD200-deficient Mice: A ROLE FOR Toll-LIKE RECEPTOR ACTIVATION. J Biol Chem. 286: 34722-32.