

Afe I

AGC[^]GCT
AccuCut™ Restriction Endonuclease

• Cat. No. E-1111 200 Units
E-1112 1000 Units

• Lot No. : 01112451H8A3

• Supplied with Enzyme

10X AccuCut™ violet Buffer : 1 mL
330mM Tris-acetate pH 7.9
100 mM Mg-acetate
660 mM K-acetate
10mM DTT

1X Dilution Buffer : 1 mL
10 mM pH 7.6 Tris-HCl
50 mM KCl
0.1 mM EDTA
1 mM DTT
200 µg/mL Acetylated BSA
50% Glycerol

• Store at -20 °C.

• **Unit definition** : One unit of restriction endonuclease activity is defined as the amount of enzyme required to completely digest 1µg of substrate DNA in a total reaction volume of 50 µL in one hour using the AccuCut™ buffer provided. Incubations are performed in 1.5 mL tubes at the appropriate incubation temperature as indicated in the Product Profile.

• **Isoschizomer** : *Ait I* ,*Aor51H I* ,*Fun I* ,*Eco47 III*.

• **Neoschizomer** : Unfound

• **Reactivity on methylated substrate DNA**:
Unidentified

• **Ref)**Abdurashitov, M.A., Kileva, E.V., Shevchenko, A.V., Degtyarev, S.K., Unpublished observations.

• **Source** :*Alcaligenes faecalis* T2774.

• **Concentration** : 20 Units/µL

• **Reaction Condition**

- 10X AccuCut™ violet Buffer
- Incubate at 37 °C.

• **Storage Buffer**

20 mM pH 7.5, Tris-HCl
50 mM KCl
1 mM EDTA
10 mM 2-mercaptoethanol
50% Glycerol

• **Heat inactivation** :65c for 20 minutes.

Quality Control

• **Overdigestion Assay** :

No nonspecific activity was detected after incubation of 1 µg of λ DNA with 50 units of *Afe I* for 15 hours.

* Conditions of low ionic strength, high enzyme concentration, glycerol concentration >5%, or pH >8.0 may result in star activity.

• **Nuclease Contamination Assay** :

No altered pattern was detected after incubation of 1 µg of substrate DNA with *Afe I* in 50 µL reaction volume with the supplied AccuCut™ buffer overnight.

• **Ligation and Recutting Assay** :

This assay is used to test for exonuclease activity that would degrade the termini of restriction fragments, resulting in inhibition of ligation and of subsequent digestion of ligated fragments. After 40-fold overdigestion with *Afe I*, 95% of the DNA fragments can be ligated and recut with *Afe I*.