

Dra I

TTT[^]AAA AccuCut™ Restriction Endonuclease

- Cat. No. E-1601 2000 Units
E-1602 10000 Units

- Lot No. : 02C151491H8A3

- Supplied with Enzyme

| | |
|----------------------------|-------------------|
| 10X AccuCut™ Orange Buffer | : 1 mL |
| 100 mM | pH 7.6 Tris-HCl |
| 100 mM | MgCl ₂ |
| 500 mM | NaCl |
| 10 mM | DTT |
| 1X Dilution Buffer | : 1 mL |
| 10 mM | pH 7.6 Tris-HCl |
| 50 mM | KCl |
| 0.1 mM | EDTA |
| 1 mM | DTT |
| 200 µg/mL | Acetylated BSA |
| 50% | Glycerol |

- Store at -20 °C.

- Source : *Deinococcus radiophilus*.

- Concentration : 40 Units/µL

- Reaction Condition

- 10X AccuCut™ Orange Buffer Buffer
- Incubate at 37 °C.

- Storage Buffer

| | |
|-----------|------------------|
| 10 mM | pH 7.5, Tris-HCl |
| 50 mM | KCl |
| 1 mM | DTT |
| 0.1 mM | EDTA |
| 0.2 mg/mL | BSA |
| 50% | Glycerol |

- Heat inactivation : 65 °C for 20 minutes.

- **Unit definition** : One unit of restriction endonuclease activity is defined as the amount of enzyme required to completely digest 1 µg of substrate DNA in a total reaction volume of 50 µL in one hour using the AccuCut™ buffer provided. Incubations are performed in 1.5 mL tubes at the appropriate incubation temperature as indicated in the Product Profile.

- **Isoschizomer** : *Aha III*.

- **Neoschizomer** : Unfound

- **Reactivity on methylated substrate DNA**: Blocked by TTTA^{m6}AA

- **Ref)** 1. Kur, J., (1993) *Acta Microbiol. Pol.*, vol. 42, pp. 145-150.
2. Morgan, R., *Unpublished observations*.
3. Purvis, I.J., Moseley, B.E.B., (1983) *Nucleic Acids Res.*, vol. 11, pp. 5467-5474.
4. Purvis, I.J., Moseley, B.E.B., (1983) *Nucleic Acids Res.*, vol. 11, pp. 5467-5474.

Quality Control

- **Overdigestion Assay** :

No nonspecific activity was detected after incubation of 1 µg of λ DNA with 50 units of *Dra I* for 15 hours.

* Conditions of low ionic strength, high enzyme concentration, glycerol concentration >5%, or pH >8.0 may result in star activity.

- **Nuclease Contamination Assay** :

No altered pattern was detected after incubation of 1 µg of substrate DNA with *Dra I* in 50 µL reaction volume with the supplied AccuCut™ buffer overnight.

- **Ligation and Recutting Assay** :

This assay is used to test for exonuclease activity that would degrade the termini of restriction fragments, resulting in inhibition of ligation and of subsequent digestion of ligated fragments. After 40-fold overdigestion with *Dra I*, 95% of the DNA fragments can be ligated and recut with *Dra I*.