

AccuPower[®] RT-PCR PreMix is covered by Korea Pat. No. 162770 and US Pat. No. 5861251 & 6153412.

Headquarters

49-3, Munpyeong-dong, Daedeok-gu, Daejeon 306-220, Korea
Phone: +82-42-936-8500
Fax: +82-42-930-8600

Seoul Office

1 floor, Donggwan Building 1566-2, Seocho-dong, Seocho-gu, Seoul 137-874 Korea
Phone: +82-2-598-1094
Fax: +82-2-598-1096

Bioneer Inc.

1000 Atlantic Avenue Alameda, CA 94501 USA
Toll free : 1-877-264-4300
Fax : 1-510-865-0350
E-mail: order.usa@bioneer.com

Order

Korea Market only: 1588-9788
E-mail: order@bioneer.co.kr
Website: www.bioneer.com

AccuPower[®] RT/PCR PreMix contains all the components necessary for cDNA synthesis and amplification in one tube.

Advantages

- Speed** Substantial reduction in reaction setup time.
- Stability** As each tube of AccuPower[®] RT/PCR PreMix contains a stabilizer (patented in US and Korea), which can maintain the stability of the RTase and DNA polymerase up to 2 years at -20°C. Pre-denaturation during PCR has no effects on the activity of the DNA polymerase.
- Reproducibility and Yield** The strict functional QC assays demonstrated highly consistent and reproducible RT/PCR performance. In most applications an increase in yield is observed as compared to the standard reactions.
- Simplicity** The fewer manual steps allow reduction in potential errors. Each tube contains tracking dye and precipitant for agarose gel electrophoresis, eliminating the needs for a separate loading buffer.

Experimental Protocol

- Mix the template RNA and the reverse primer in a sterile tube as indicated below;

Concentration of template RNA and primer			
Reaction volume		20 µl reaction	50 µl reaction
Template RNA	Total RNA	0.5~1.0 µg	1.0~2.0 µg
	Poly(A) RNA	0.05~0.1 µg	0.1~0.2 µg
Primer	Primer	10~30 pmole	20~50 pmole

- Incubate the mixture at 70°C for 5min and place it on ice.
- Transfer the incubated mixture and the forward primer to AccuPower[®] RT/PCR PreMix tube, then fill up the reaction volume with DEPC-DW.
- Dissolve the lyophilized blue pellet by vortexing, and briefly spin down.
- Add mineral oil to each tube (This step is unnecessary when using a thermal cycler with top heating.)
- Perform cDNA synthesis reaction as follows:
 - 42°C, 60 min. (cDNA synthesis)
 - 94°C, 5 min. (RTase inactivation)
- Perform PCR cycles according to the PCR condition.
(Annealing temperature and time need to be optimized for each primer/template combination.)

Trademark AccuPower is a trademark of Bioneer Corporation.

Ordering Information

Tube type	Reaction	Cat. No	Description	Tube type	Reaction	Cat. No	Description
0.2ml Tube	20 ul	K-2055	0.2 ml thin-wall 8-strip tubes with attached cap / 96 tubes	96-well	20 ul	K-2262-4	thin-wall 96-well flat plate
		K-2055-B	0.2 ml thin-wall 8-strip tubes with attached cap / 480 tubes			K-2262-5	thin-wall 96-well full-skirted plate
	50 ul	K-2057	0.2 ml thin-wall 8-strip tubes with attached cap / 96 tubes			K-2262-6	thin-wall 96-well semi-skirted plate
		K-2057-B	0.2 ml thin-wall 8-strip tubes with attached cap / 480 tubes			K-2084-1	thin-wall 384-well full-skirted plate
96-well	10 ul	K-2262-1	thin-wall 96-well flat plate	384-well	10 ul	K-2084-2	thin-wall 384-well full-skirted plate
		K-2262-2	thin-wall 96-well full-skirted plate		20 ul	K-2084-3	thin-wall 384-well full-skirted plate
		K-2262-3	thin-wall 96-well semi-skirted plate	0.5ml Tube	50ul	k-2056	0.5 ml thin-wall tubes with attached cap / 100 tubes

A complete product list appears on our web site at www.bioneer.com