MONOCLONAL ANTIBODY



Anti-Peroxiredoxin II (1E8)

Background : Peroxiredoxin (Prx) is a growing peroxidase family, whose mammalian members have been known to connect with cell proliferation, differentiation, and apoptosis.

Many isoforms (about 50 proteins), collected in accordance to the amino acid sequence particularly amino-terminal homology, region containing active site cysteine residue, and the thiol-specific antioxidant activity, distribute throughout all the kingdoms. Among them, mammalian Prx consists of 6 different members grouped into typical 2-Cys, atypical 2-Cys Prx, and 1-Cys Prx. Except Prx VI belonging to 1-Cys Prx subgroup, the other five 2-Cys Prx isotypes have the thioredoxin-dependent peroxidase utilizing activity thioredoxin, thioredoxin reductase, and NADPH as a reducing system. Mammalian Prxs are 20 -30 kilodalton in molecular size and vary in subcellular localization: Prx I, II, and VI in cytosol, Prx III in mitochondria, Prx IV in ER and secretion, Prx V showing complicated including distribution peroxisome, mitochondria and cytosol (1).

Immunogen: Recombinant human protein

purified from *E.coli*

Host: Mouse

Clone number: 1E8

Isotype: IgG1, ĸ

Size: 100 μℓ

Composition: PBS containing 50% glycerol


Positive control: HeLa cell lysate

Storage : Store for 1 year at -20°C from date

of shipment

Species cross reactivity

Human	Mouse	Rat
+	+	+



IMMUNOBLOT ANALYSIS of cell lysates

Lane 1 : HeLa cell lysate Lane 2 : 293T cell lysate Lane 3 : SH-SY5Y cell lysate Lane 4 : HepG2 cell lysate

Applications:

ELISA

Western Blotting(1:2,000~4,000)

Immunohistochemistry

Background Reference:

1) Wood, Z. A. et al. (2003) Trends Biochem Sci.

28(1):32-40.

FOR RESEARCH PURPOSE ONLY NOT FOR DIAGNOSTIC OR THERAPEUTIC USE