

Protein Phosphatase 2A catal. subunit-alpha/beta Human, Mouse Monoclonal Antibody, Clone: 20G6

Product Data Sheet Cat. No.:

Source of Antigen: Synthetic peptide (a.a. 297- 309) LF-MA0220 (100 ul)

Host: Mouse

Isotype: IgG2b, kappa

Other names: PP2A

Preparation

Hepes with 0.15M NaCl, 0.01% BSA, 0.03% sodium azide, and 50% glycerol

Species Reactivity

Human (+), Mouse (+), Rat (+)

Storage/Stability

Store for 1 year at -20°C from date of shipment

Applications

Immunoprecipitation, Western blotting

Introduction to the Molecule

Protein Phosphatase 2A (PP2A) is a ubiquitous and conserved Serine/Threonine phosphatase and accounts for a large fraction of phosphatase activity in eukaryotic cells. PP2A plays an important role in cell cycle regulation, cell growth control, development, egulation of various signal transduction pathways, and cell mobility. PP2A comprises A and B subunits which are regulatory and a catalytic C subunit. When the PP2A catalytic C subunit (36 kDa) associates with the regulatory A (65 kDa, PR65) and B subunits (PR55, PR56, PR72, PR93 etc), wide variety of heterotrimeric holoenzymes are produced with distinct functions and characteristics. The different association of the subunits give PP2A large regulatory flexibility and differential substrate specificity. The A subunit exists as two isoforms (alpha and beta) as does the C subunit, whereas the B subunits fall into three families designated B, B' (also called B56), and B''. The A subunit is the scaffold required for the formation of the heterotrimeric complex and the binding of A subunit alters the enzymatic activity of the catalytic subunit, even if the B subunit is absent. Storage: Store for 1 year at -20°C from date of shipment Applications

Gentaur Molecular Products Voortstraat 49 1910 Kampenhout, Belgium http://www.gentaur-worldwide.com