MONOCLONAL ANTIBODY



Anti-Casein kinase II α (27H4)

Background : The casein kinase I (CKI) family of serine/threonine protein kinases is highly conserved from yeast to humans. The CKI family is involved in many diverse and important cellular functions, such regulation of membrane transport, cell division, DNA repair, circadian rhythms, and nuclear localization.

The name of the enzyme family was originated from the convenience of casein as a substrate since the earliest days of research on protein phosphorylation.

Casein kinase 2 (CK2) is a ubiquitous, highly conserved, essential serine/threonine kinase which has been implicated in cell cycle control, DNA repair, regulation of the circadian rhythm and other cellular processes.

CK2 is a tetramer of two alpha subunits and two beta subunits. The alpha subunits have the catalytic kinase domain. Loss of the α' catalytic subunit produces male infertility, and loss of the single β regulatory subunit is early embryonic lethal.

CK2 appears to be upregulated in most cancers and the promotion of tumorigenesis by the overexpression of CK2 has been reported in transgenic mice.

Immunogen: Recombinant human protein purified from *E.coli* (His-Casein kinase II a)

Host: Mouse

Clone number: 27H4

Isotype : IgG2b, k

Size: 100ul

Western Blotting (1:2,000)

Applications:

ELISA


Background Reference:

- 1) Price M.A., 2006 Genes Dev. 20:399-410
- 2) Seldin DC et al., 2005, Mol Cell Biochem. 274(1-2):63-67.
- 3) Tobin A.B., 2002, Trends Pharmacol Sci. 23:337-343
- 4) Gross S.D. and Anderson R.A., 1998, Cell Signal. 10:699-711

Composition: Hepes with 0.15M NaCl, 0.01% BSA, 0.03% sodium azide, and 50% glycerol **Positive control:** Jurkat T cell lysate

Storage : Store for 1 year at -20°C from date of shipment

Species cross reactivity Human Mouse Rat +



FOR RESEARCH PURPOSE ONLY NOT FOR DIAGNOSTIC OR THERAPEUTIC USE