



Cartilage Oligomeric Matrix Protein Human HEK293

Product Data Sheet

Type: Recombinant

Source: HEK293

Species: Human

Other names: COMP, Thrombospondin-5, TSP5

Cat. No.:

RD172080100 (0.1 mg)

Description

Total 750 AA. MW: 82.4 kDa (calculated). N-Terminal FLAG-tag, 13 extra AA (highlighted). Protein identity confirmed by LC-MS/MS.

Introduction to the Molecule

Cartilage oligomeric matrix protein (COMP), also designated thrombospondin 5 (TSP 5), is non-collagenous glycoprotein and is a member of the thrombospondin family of extracellular proteins. COMP is a calcium-binding protein of high molecular weight (>500kDa) present in the extracellular matrix of articular, nasal and tracheal cartilage. COMP is not only cartilage-derived but was found widely in other tissues, including synovium and tendon. Intact COMP is pentameric, with five identical subunits and the carboxy-terminal globular domain of native COMP binds to collagens I, II, and IX. It has been proposed that COMP molecules are important for maintaining the properties and integrity of collagen network. In addition COMP may have a storage and delivery function for hydrophobic cellsignaling molecules such as vitamin D. The significance of COMP for normal development and function of cartilage has been underscored by the discovery that mutations of the COMP gene result in pseudoachondro-plasia and some forms of multiple epiphyseal dysplasia. Most published studies have shown that serum levels of COMP provide important information about metabolic changes occurring in the cartilage matrix in joint disease. These studies describe that serum COMP level correlated with cartilage degradation and is a potential prognostic marker in inflammatory joint diseases such as osteoarthritis (OA) and rheumatoid arthritis (RA). Results have demonstrated an association of increasing serum COMP levels with progressive destruction of articular cartilage monitored radiographically. OA and RA are a common disease causing pain and disability in a significant proportion of the adult population and early diagnostics of these diseases is very important for future therapy.

Research topic

Bone and cartilage metabolism

Amino Acid Sequence

HVDYKDDDDK **PAG**QQQSPLG SDLGPMMLRE LQETNAALQD VRELLRQQVR EITFLKNTVM ECDACGMQQS VRTGLPSVRP
LLHCAPGFCE PGVACIQTES GARGPCPCAG FTGNNGSHCTD VNECNAHPCF PRVRCINTSP GFRCEACPPG YSGPTHQGVG
LAFKANKQV CTDINECETG QHNCVPNSVC INTRGSFQCG PCQPGFVGDQ ASGCQRRQR FCPDGPSPSEC HEHADCVLER
DGSRSVCVAV GWAGNGILCG RDTDLGDFPD EKLRCPERQC RKDNCVTVPN SGQEDVDRDG IGDACDPDAD GDGVPNEKDN
CPLVRNPQQR NTDEDKWGDA CDNCRSQKND DQKDTDQDGR GDACDDDDIDG DRIRNQADNC PRVPNSDQKD SDGDGIGDAC
DNCPQKSNPD QADVHDHDFVG DACDSDDQDQD GDGHQDSRDN CPTVPNSAQE DSDHDGQGDG CDDDDDDNDGV PDSRDNCRLV
PNPGQEDADR DVGVDVCQDD FDADKVVDKI DVCPEAEVT LTDFRAFQTV VLDPEGDAQI DPNWVVLNQG REIVQTMNSD
PGLAVGYTAF NGVDFEGTFH VNTVTDDDYA GFIFGYQDSS SFYVVMWKQM EQTYWQANPF RAVAEPGIQL KAVKSSTGPG
EQLRNALWHT GDTEQVRLR WKDPRNVGWK DKKSRYWFLQ HRPQVGYIRV RFYEGPELVA DSNVVLDTTM RGGRLGVFCF
SQENIIWANL RYRCNDTIPE DYETHQLRQA


Source

HEK293

Purity

Purity as determined by densitometric image analysis: >90%

SDS-PAGE gel



10% SDS-PAGE separation of Human COMP

1. M.W. marker - 21, 31, 45, 66, 97 kDa

2. reduced and boiled sample, 5µg / lane

3. non-reduced and non-boiled sample, 5µg / lane

Endotoxin

< 1.0 EU/ug

Formulation

Filtered (0.4 µm) and lyophilized from 0.5 mg/ml in 20mM Tris buffer, 50mM NaCl, pH 7.5

Reconstitution

Add deionized water to prepare a working stock solution of approximately 0.5 mg/mL and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.

Shipping

At ambient temperature. Upon receipt, store the product at the temperature recommended below.

Storage, Stability/Shelf Life

Store lyophilized protein at -20°C. Lyophilized protein remains stable until the expiry date when stored at -20°C. Aliquot reconstituted protein to avoid repeated freezing/thawing cycles and store at -80°C for long term storage. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after one week at 4°C.

Quality Control Test

BCA to determine quantity of the protein.

SDS PAGE to determine purity of the protein.

LAL to determine quantity of endotoxin.

Applications

Cell culture and/or animal studies, ELISA, Western blotting

Note

This product is intended for research use only. The Certificate of Analysis is available on www.biovendor.com

Physical Appearance

Filtered (0.4 µm) white lyophilized (freeze-dried) powder.

Gentaur Molecular Products
Voortstraat 49
1910 Kampenhout, Belgium
<http://www.gentaur-worldwide.com>