



## **Cartilage Oligomeric Matrix Protein (COMP) Human, Mouse Monoclonal Antibody, Clone: 12C4**

### **Product Data Sheet**

**Source of Antigen:** Human articular cartilage

**Host:** Mouse

**Isotype:** IgG1

**Cat. No.:**

RD182080100C4 (0.1 mg)

**Other names:** COMP, Thrombospondin-5, TSP5

### **Research topic**

Bone and cartilage metabolism

### **Preparation**

The antibody is a mouse monoclonal antibody against Human COMP.

### **Purification Method**

Affinity chromatography on a column with immobilized protein G.

### **Antibody Content**

0.1 mg (determined by BCA method)

### **Formulation**

The antibody is lyophilized in 0.05 M phosphate buffer, 0.1 M NaCl, pH 7.2. **AZIDE FREE.**

### **Reconstitution**

Add 0.1 ml of deionized water and let the lyophilized pellet dissolve completely. Slight turbidity may occur after reconstitution, which does not affect activity of the antibody. In this case clarify the solution by centrifugation.

### **Shipping**

At ambient temperature. Upon receipt, store the product at the temperature recommended below.

### **Storage/Stability**

The lyophilized antibody remains stable and fully active until the expiry date when stored at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles and store frozen at -80°C. Reconstituted antibody can be stored at 4°C for a limited period of time; it does not show decline in activity after one week at 4°C.

### **Expiration**

See vial label.

### **Lot Number**

See vial label.

### **Quality Control Test**

Indirect ELISA - to determine titer of the antibody

SDS PAGE - to determine purity of the antibody

### **Applications**

ELISA, Immunohistochemistry

## Introduction to the Molecule

Cartilage oligomeric matrix protein (COMP), also designated thrombospondin 5 (TSP 5), is non-collagenous glycoprotein and is a member of the thrombospondin family of extracellular proteins. COMP is a calcium-binding protein of high molecular weight (>500kDa) present in the extracellular matrix of articular, nasal and tracheal cartilage. COMP is not only cartilage-derived but was found widely in other tissues, including synovium and tendon. Intact COMP is pentameric, with five identical subunits and the carboxy-terminal globular domain of native COMP binds to collagens I, II, and IX. It has been proposed that COMP molecules are important for maintaining the properties and integrity of collagen network. In addition COMP may have a storage and delivery function for hydrophobic cellsignaling molecules such as vitamin D. The significance of COMP for normal development and function of cartilage has been underscored by the discovery that mutations of the COMP gene result in pseudoachondro-plasia and some forms of multiple epiphyseal dysplasia. Most published studies have shown that serum levels of COMP provide important information about metabolic changes occurring in the cartilage matrix in joint disease. These studies describe that serum COMP level correlated with cartilage degradation and is a potential prognostic marker in inflammatory joint diseases such as osteoarthritis (OA) and rheumatoid arthritis (RA). Results have demonstrated an association of increasing serum COMP levels with progressive destruction of articular cartilage monitored radiographically. OA and RA are a common disease causing pain and disability in a significant proportion of the adult population and early diagnostics of these diseases is very important for future therapy.

## References to this Product

- Kraus VB, Huebner JL, Stabler T, Flahiff CM, Setton LA, Fink C, Vilim V, Clark AG. *Ascorbic acid increases the severity of spontaneous knee osteoarthritis in a guinea pig model.* Arthritis Rheum. 2004 Jun;50 (6):1822-31
- Misumi K, Vilim V, Clegg PD, Thompson CC, Carter SD. *Measurement of cartilage oligomeric matrix protein (COMP) in normal and diseased equine synovial fluids.* Osteoarthritis Cartilage. 2001 Feb;9 (2):119-27
- Vilim V, Voburka Z, Vytasek R, Senolt L, Tchetverikov I, Kraus VB, Pavelka K. *Monoclonal antibodies to human cartilage oligomeric matrix protein: epitope mapping and characterization of sandwich ELISA.* Clin Chim Acta. 2003 Feb;328 (1-2):59-69
- Misumi K, Vilim V, Hatazoe T, Murata T, Fujiki M, Oka T, Sakamoto H, Carter SD. *Serum level of cartilage oligomeric matrix protein (COMP) in equine osteoarthritis.* Equine Vet J. 2002 Sep;34 (6):602-8

## Note

This product is for research use only.

**Gentaur Molecular Products**  
**Voortstraat 49**  
**1910 Kampenhout, Belgium**  
**<http://www.gentaur-worldwide.com>**