www.biossusa.com support@biossusa.com 800.501.7654 [DOMESTIC] +1.781.569.5821 [INTERNATIONAL]

# Bioss

# bs-0521R-PE-Cy3

· Rabbit Anti-CD44 Polyclonal Antibody, PE-Cy3 conjugated

Conjugated Primary Antibodies

#### Background:

The protein encoded by this gene is a cell-surface glycoprotein involved in cell-cell interactions, cell adhesion and migration. It is a receptor for hyaluronic acid (HA) and can also interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs). This protein participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing, hematopoiesis, and tumor metastasis. Transcripts for this gene undergo complex alternative splicing that results in many functionally distinct isoforms, however, the full length nature of some of these variants has not been determined. Alternative splicing is the basis for the structural and functional diversity of this protein, and may be related to tumor metastasis. [provided by RefSeq, Jul 2008].

Purification: Was purified by Protein A and peptide affinity chromatography.

### Storage:

Aqueous buffered solution containing 100ug/ml BSA, 50% glycerol and less than 0.09% sodium azide. Store at -20°C for 12 months. Protect from light. [Product without BSA and/or sodium azide is available for special order.]

#### Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

For full size images and description please click  $\ensuremath{\mathsf{HERE}}\,.$ 

Size: 100ul

Concentration: 1ug/uL

Host: Rabbit Reactivities:

Human, Mouse, Rat, Dog, Pig, Bovine, Rabbit,

## Application:

- IF(1:100-500)
- Not yet tested in other applications.
  Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 85kDa

Note:

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.