

bs-1322R-PE

• Rabbit Anti-gamma tubulin(Centrosome Marker) Polyclonal Antibody, PE conjugated

Conjugated Primary Antibodies

Background:

Centrosome Marker

Gamma tubulin, a member of the tubulin superfamily, is a ubiquitous and highly conserved protein within the microtubule organizing centre (MTOC). Gamma tubulin is not a component of microtubules, rather it functions as the microtubule nucleator at the MTOC, is responsible for binding microtubule minus ends and mediating the link between microtubules and the centrosome. By binding to the beta tubulin subunit of the tubulin molecule, it establishes the polarity of a microtubule leaving the alpha tubulin subunit exposed at the positive end. The abundance of Gamma tubulin is less than 1% of the level of either alpha or beta tubulin. It shares approximately 28-32% identity with alpha tubulin from various organisms and 32-36% identity with beta tubulins. The detection, localization and characterization of proteins involved in microtubule function is fundamental to the understanding of mitosis, meiosis and the microtubule cytoskeleton.

Purification: Was purified by Protein A and peptide affinity chromatography.


Storage:

Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year. Protect from light.

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.



For full size images and description please click [HERE](#).

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities:

Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse, Rabbit, Guinea Pig,

Application:

- IF(1:100-500)
- Not yet tested in other applications. Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 50kDa

Preservatives: 10ug/uL BSA and 0.1% NaN₃.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.