bs-1796R-FITC

· Rabbit Anti-GCK/Glucokinase Polyclonal Antibody, FITC conjugated

Conjugated Primary Antibodies

Background:

Hexokinases phosphorylate glucose to produce glucose 6 phosphate, thus committing glucose to the glycolytic pathway. Alternative splicing of this gene results in three tissue specific forms of glucokinase, one found in pancreatic islet beta cells and two found in liver. The protein localizes to the outer membrane of mitochondria. In contrast to other forms of hexokinase, this enzyme is not inhibited by its product glucose 6 phosphate but remains active while glucose is abundant. Mutations in this gene have been associated with non insulin dependent diabetes mellitus, also called maturity onset diabetes of the young, type 2; mutations have also been associated with persistent hyperinsulinemic hypoglycemia of infancy (PHHI).

Purification: Was purified by Protein A and peptide affinity chromatography.

Storage:

Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year. Protect from light.

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities:

Human, Mouse, Rat, Chicken, Dog, Pig, Horse, Rabbit,

Application:

IF(1:100-500)

 Not yet tested in other applications.
Optimal working dilutions must be determined by the end user.

Antibody Type: Polydonal

Isotype: IgG

Molecular Weight: 51kDa

Preservatives:

10ug/uL BSA and 0.1% NaN3.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.

