

www.biossusa.com
support@biossusa.com
800.501.7654 [DOMESTIC]
+1.781.569.5821 [INTERNATIONAL]



bs-9882R-Gold

• Rabbit Anti-GLCNE Polyclonal Antibody, Gold conjugated

Conjugated Primary Antibodies

Background:

The bifunctional enzyme UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase (GNE/Mnk), or GLCNE, regulates and initiates biosynthesis of N-acetylneuraminic acid (NeuAc), a precursor of sialic acids. GLCNE is required for normal sialylation in hematopoietic cells. Sialylation is implicated in cell adhesion, signal transduction, tumorigenicity and metastatic behavior of malignant cells. It is upregulated after PKC-dependent phosphorylation and is most abundantly expressed in liver and placenta. It is also expressed, to a lesser extent, in heart, brain, lung, kidney, skeletal muscle and pancreas. Defects in GLCNE are the cause of sialuria, inclusion body myopathy 2 (IBM2) and Nonaka myopathy (NM) or distal myopathy with rimmed vacuoles (DMRV). Sialuria is an autosomal dominant disorder caused by a lack of feedback inhibition of GLCNE by CMP-NeuAc, resulting in overproduction of NeuAc. It is characterized by an accumulation of free sialic acid in the cytoplasm and large quantities of neuraminic acid in the urine. Both IBM2 and NM/DMRV are autosomal recessive neuromuscular disorders characterized by adult onset, distal and proximal muscle weakness (especially in the legs) and a typical muscle pathology including filamentous inclusions and rimmed vacuoles.

Purification: Was purified by Protein A and peptide affinity chromatography.

Storage: Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year.

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities:

Human, Mouse, Rat, Pig, Cow, Horse, Rabbit, Sheep,

Application:

- Electron Microscopy
- Not yet tested in other applications. Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 79kDa

Preservatives:

10ug/uL BSA and 0.1% NaN3.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.

For full size images and description please click [HERE](#).

