

www.biossusa.com  
 support@biossusa.com  
 800.501.7654 [DOMESTIC]  
 +1.781.569.5821 [INTERNATIONAL]



## bs-9895R

### • Rabbit Anti-CNO/Cappuccino Polyclonal Antibody

#### Primary Antibodies

#### Background:

Biogenesis of lysosome-related organelles complex-1 (BLOC-1) is a multisubunit protein necessary for biogenesis of specialized organelles of the endosomal-lysosomal system (such as melanosomes and platelet-dense granules). The complex consists of coiled-coil-forming proteins Snapin, Pallidin, Cappuccino, Muted, BLOS1, BLOS2, and BLOS3. The localization of these proteins varies as they can be cytoplasmic, peripheral membrane bound or anchored to the vesicular membrane. Cappuccino, a primarily cytoplasmic protein, plays a role in the development of melanosomes, platelet-dense granules and other lysosome-related organelles. It interacts primarily with pallidin and Muted and has been implicated as an actin-nucleation factor that may play a role in crosstalk between microfilaments and microtubules.

#### Source/Purification:

KLH conjugated synthetic peptide derived from human CNO/Cappuccino. Was purified by Protein A and peptide affinity chromatography.

**Storage:** Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year.

#### Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

For full size images and description please click [HERE](#).

**Size:** 100ul or 100ug lyophilized

**Concentration:** 1ug/uL

**Host:** Rabbit

#### Reactivities:

Human, Mouse, Rat, Dog, Pig, Cow, Horse, Sheep, Guinea Pig, hu, mo, rat, pig, hrs, cow, shp, dog, Gpig

#### Application:

- WB(1:100-500)
- ELISA(1:500-1000)
- IHC-P(1:100-500)
- IHC-F(1:100-500)
- IF(1:50-200)
- Not yet tested in other applications. Optimal working dilutions must be determined by the end user.

**Antibody Type:** Polyclonal

**Isotype:** IgG

**Molecular Weight:** 27kDa

**Preservatives:** 10ug/uL BSA and 0.1% NaN<sub>3</sub>.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.

