

Rat anti mouse CD126 (IL-6R α chain)

(Azide-free & Low endotoxin)

ORDERING INFORMATION

Catalog Number: gAP-0041 Size: 1.00 mg Storage: <-20° C

Immunogen: Mouse recombinant IL-6R α chain

Ig Type: Rat IgG2
Clone AP-MAB0848
Endotoxin Level < 0.002EU/µg IgG*
Applications: FC, IP & Blocking

Description: CD126 is an 80 kD IL-6 receptor α chain also known as IL-6R. It is a member of the immunoglobulin superfamily that is expressed on activated T and B cells, monocytes, hepatocytes, and plasma cells. High affinity IL-6 receptors are formed by the non-covalent association of CD126 and the IL-6 receptor β chain (CD130 or gp130). CD126 binds IL-6 with low affinity, but does not signal. The β chain (gp130, CD130) does not bind IL-6 by itself, but associates with the α -chain/IL-6 complex to initiate signal transduction. IL-6 binding to the receptor complex results in the stimulation of B and T cells, and hematopoietic precursor stimulation and differentiation.

Preparation: This antibody was produced from a hybridoma (mouse myeloma fused with spleen cells from a rat immunized with **Mouse recombinant IL-6R** α **chain**.

Formulation: The IgG fraction of culture supernatant was purified by Protein A/G affinity chromatography and lyophilized from a 0.2 μm filtered solution in phosphate-buffered saline (PBS, Azide Free).

Reconstitution: Reconstitute the antibody with sterile PBS and the reconstituted antibody can be aliquoted and stored frozen at < -20 for at least for six months without detectable loss of activity. **Avoid repeated freeze-thaw cycles.** Lyophilized samples are stable for 2 years from date of receipt when stored at -70°C.

*Endotoxin Level: Extremely low level of LPS (< 0.002EU/µg IgG)

Application(s):

- 1. FC and IP
- 2. Blocking IL-6/IL-6R interactions

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Tel: 617-549-2665; Fax: (480) 247-4337, angioproteomie@gmail.com

^{*} The antibody is produced by in vitro culture.