

## Rabbit anti EGFR (pTyr1197)

Catalog Number: pAP11228

Amount: 100µl (for 10 Western Blotting)

Swiss-Prot No.: P00533

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150mM NaCl,

0.02% sodium azide and 50% glycerol.

Storage/Stability: Store at -20 °C/1 year

Immunogen: The antiserum was produced against synthesized phosphopeptide derived from human

EGFR around the phosphorylation site of tyrosine 1197 (A-E-Y<sup>P</sup>-L-R).

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatogramphy using non-phosphopeptide corresponding to the phosphorylation site.

Specificity/Sensitivity: EGFR (phospho-Tyr1197) antibody detects endogenous levels of EGFR only when

phosphorylated at tyrosine 1197

Reactivity: Human, Mouse, Rat


Applications: WB: 1:1000

References: Corbalan-Garcia S, et al. (1996) Mol Cell Biolt; 16(10): 5674-5682

Kanner SB, et al. (1991) Mol Cell Biol; 11(2): 713-720 Wu TT, et al. (1998) Mol Biol Cell; 9(7): 1661-1674

O'Rourke DM, et al. (1997) Proc Natl Acad Sci U S A; 94(7): 3250-3255

Goldstein DJ, et al. (1994) J Virol; 68(7): 4432-4441



Western blot analysis of extract from A431 cells untreated or treated with EGF (200ng/ml, 5min), using EGFR (Ab-1197) antibody (AP21221, Lane 1 and 2) and EGFR (phospho-Tyr1197) antibody (pAP11228, Lane 3 and 4).

## For Research Use Only

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Fax: (480) 247-4337, angioproteomie@gmail.com