

Biotinylated Anti-mouse CD105/Endoglin (MJ7/18)

Description: The MJ7/18 monoclonal antibody reacts with the mouse CD105 molecule, also known as Endoglin. This 90 kDa disulfide-linked homodimer is expressed by vascular endothelial cells. It is suggested that CD105 functions in adhesion and embryonic angiogenesis. The rat antibody was purified from cell culture supernatant by Protein G chromatography and then biotinylated using a standard protocol.

Host species Rat

Antigen: Inflamed mouse skin
Purification: Protein G chromatography

Stabilizer: none

Buffer: PBS pH 7.4 w/o preservative

Formulation: lyophilized rat IgG2a

Reconstitution: The biotinylated antibody should be reconstituted to a concentration of 50 μ g/ml with sterile PBS solution containing 0.1% BSA. This solution can be stored at 4°C for at least one month without detectable loss of activity. Frozen aliquots of this solution are stable for at least 6 months when kept at -20°C. Avoid more than one freeze-thaw cycle.

Specificity: The unconjugated antibody can be used for ELISA experiments, Western blotting, immunoprecipitation, FACS and cell sorting.

Applications

Western Analysis: Use at a concentration of 1-2 µg/ml with the appropriate secondary reagents.

Immuoprecipitation: Use at a concentration of 1-2 μ g with the appropriate secondary reagents.

FACS analysis: Use at 3-20 µg/ml together with the appropriate secondary reagents

Immunohistochemistry: It was reported for use of staining of frozen sections.

Optimal dilutions should be determined by each laboratory for each application.

Usage: Anti-mouse CD105/Endoglin is offered for research use. Not for drug use. Not for human use.

Catalogue number: 103-MBi60 Size: 50 μg

Literature: [Ge, AZ and Butcher EC, Gene 138:201-206, 1994]

** please note: always centrifuge vials before opening **