

RayBiotech, Inc.

3607 Parkway Lane suite 200 Norcross,GA 30092 Tel: 770-729-2992, 1-888-494-8555

Fax: 770-206-2393

Website: www.raybiotech.com Email: info@raybiotech.com

Certificate of Analysis and Data Sheet

Recombinant Mouse Interferon-gamma

Catalog No.
Source:
Escherichia Coli

Synonyms

Immune Interferon, type II interferon, T cell interferon, MAF, IFNG, IFG, IFI, IFN-gamma.

Introduction

IFN-gamma produced by lymphocytes activated by specific antigens or mitogens.

IFN-gamma, in addition to having antiviral activity, has important immunoregulatory functions, it is a potent activator of macrophages, and has antiproliferative effects on transformed cells and it can potentiate the antiviral and antitumor effects of the type I interferons.

Description

Interferon-gamma Mouse Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 134 amino acids and having a molecular mass of 15601 Dalton. The IFN-gamma is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation

Lyophilized from a 0.2µm filtered concentrated (1mg/ml) solution in PBS, pH 7.4.

Solubility

It is recommended to reconstitute the lyophilized Interferon-gamma in sterile $18M\Omega$ -cm H_2O not less than $100\mu g/ml$, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized Interferon-gamma although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IFN-gamma should be stored at 4°C between 2-7 days and for future use below -18°C.

For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

Please prevent freeze-thaw cycles.



RayBiotech, Inc.

3607 Parkway Lane suite 200 Norcross,GA 30092 Tel: 770-729-2992, 1-888-494-8555

Fax: 770-206-2393

Website: www.raybiotech.com Email: info@raybiotech.com

Purity

Greater than 95.0% as determined by

- (a) Analysis by RP-HPLC.
- (b) Analysis by SDS-PAGE.

Amino acid sequence

The sequence of the first five N-terminal amino acids was determined and was found to be Met-His-Gly-Thr-Val.

Biological Activity

The specific activity as determined by the cytopathic affect inhibition assay with murine NIH 3T3 cells chalenged with EMC virus was < 0.1 ng/ml, corresponding to a specific activity of 10MU/mg.