



RayBiotech, Inc.

3607 Parkway Lane suite 200
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

Certificate of Analysis and Data Sheet Recombinant Mouse Interleukin-2 (IL-2)

Catalog No.
228-10902

Source:
Escherichia Coli.

Synonyms

Interleukin-2, T-cell growth factor (TCGF), Aldesleukin, Lymphokine, IL-2.

Introduction

IL2 is a secreted cytokine that is important for the proliferation of T and B lymphocytes. The receptor of this cytokine is a heterotrimeric protein complex whose gamma chain is also shared by interleukin 4 (IL4) and interleukin 7 (IL7). The expression of this gene in mature thymocytes is mono-allelic, which represents an unusual regulatory mode for controlling the precise expression of a single gene. The targeted disruption of a similar gene in mice leads to ulcerative colitis-like disease, which suggests an essential role of this gene in the immune response to antigenic stimuli.

Description

Interleukin-2 Mouse Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 148 amino acids and having a molecular mass of 17231 Dalton. The IL-2 is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation

The protein (1mg/ml) was lyophilized extensive dialysis against 10mM sodium citrate buffer to a pH=4.

Solubility

It is recommended to reconstitute the lyophilized Mouse IL-2 in sterile 18MΩ-cm H₂O not less than 100μg/ml, which can then be further diluted to other aqueous solutions.


Stability

Lyophilized Interleukin-2 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL2 should be stored at 4°C between 2-7 days and for future use below -18°C.

For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

Please prevent freeze-thaw cycles.

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**



RayBiotech, Inc.

3607 Parkway Lane suite 200
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

Purity

Greater than 98.0% as determined by

- (a) Analysis by RP-HPLC.
- (b) Analysis by SDS-PAGE.

Amino acid sequence

PTSSSTSSST AEAQQQQQQQ QQQQHHLEQL LMDLQELLSR MENYRNLLKP RMLTFKFYLP
KQATELKDLQ CLEDELGPLR HVLDTQSKS FQLEDAENFI SNIRVTIVKL KGSDNTFECQ
FDDESATVVD FLRRWIAFCQ SIISTSPQ.

Biological Activity

The ED50 as determined by the dose-dependant stimulation of murine CTLL-2 cells is 0.18 ng/ml corresponding to a Specific Activity of 55MIU/mg.

Protein content

Protein quantitation was carried out by two independent methods

1. UV spectroscopy at 280 nm using the absorbency value of 0.482 as the extinction coefficient for a 0.1% (1mg/ml) solution. This value is calculated by the PC GENE computer analysis program of protein sequences (IntelliGenetics).
2. Analysis by RP-HPLC, using a calibrated solution of IL-2 as a Reference Standard.

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**