

RayBiotech, Inc.

3607 Parkway Lane suite 200 Norcross,GA 30092 Tel: 770-729-2992, 1-888-494-8555 Fax: 770-206-2393

Website: www.raybiotech.com Email: info@raybiotech.com

Certificate of Analysis and Data Sheet

MOUSE ANTI HUMAN AQUAPORIN 4

With HRP-conjugated Secondary Antibody

Catalog No.	Species	Isotype	
DS-MB-00095	Human	IgG1	

Preparation

Preparation: Purified IgG prepared by affinity chromatography on Protein G from tissue culture

supernatant.

Immunogen: Synthetic peptide corresponding to amino acids 301-318 of rat aquaporin 4.

Fusion Partners: cells from immunized Balb/c mice were fused with the cells of Sp-2/0 Ag14 mouse

myeloma cell line.

Specificity

This product recognizes an epitope within the cytoplasmic domain of the water-specific channel aquaporin 4 (also known as AQP4). Aquaporin 4 is located in various tissues including the gastro intestinal tract, kidney and brain.

Species Cross Reactivity: Does not react with Rat, Mouse

Formulation

Product Type: Monoclonal Antibody **Product Form**: Purified IgG - liquid

Preservative Stabilizers: 0.09% Sodium Azide, 0.1% Bovine Serum Albumin.

Approx. Protein Concentrations: IgG concentration 1.0mg/ml.

Storage

18 months from date of dispatch.

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted. Storage in frost-free freezers is not recommended. Should this product contain a precipitate we recommend microcentrifugation before use.

Avoid repeated freezing and thawing as this may denature the antibody.



RayBiotech, Inc.

3607 Parkway Lane suite 200 Norcross,GA 30092 Tel: 770-729-2992, 1-888-494-8555

Fax: 770-206-2393 Website: www.raybiotech

Website: www.raybiotech.com Email: info@raybiotech.com

Applications

Table Summary of antibody applications and working conditions

Options Functions	YES	NO	Not determined	Recommended Work dilution or concentration
ELISA			•	
Western Blotting			•	
Immunohistology – frozen	•			1:100
Immunohistology – paraffin(*)	•			1:100
Immunohistology - resin				
Immunoprecipitation				
Flow Cytometry			-	
Immunodiffusion				
Neutralization				

Note: Other applications are not tested yet. Optimal dilutions should be determined by each laboratory for each application.

(*)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH6.0 is recommended for this purpose. Histology Positive Control Tissue Kidney

Secondary Antibody Applications

Options Functions	YES	NO	Not determined	Recommended Work dilution or concentration
Immunoassay (ELISA, Western blot)	•			1:5000-1:10000

Reference

Bodis, B. et al. (2001) Active water selective channels in the stomach: investigation of aquaporins after ethanol and capsaicin treatment in rats. J. Physiol. 95: 271 - 275.