



RayBiotech, Inc.

3607 Parkway Lane suite 200
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

Certificate of Analysis and Data Sheet

MOUSE ANTI RAT CD31

With HRP-conjugated Secondary Antibody

| Catalog No. | Species | Isotype |
|-------------|---------|------------------|
| DS-MB-00600 | Rat | IgG ₁ |

Preparation

Preparation: Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Immunogen: Activated, Lewis rat derived microglial cells

Fusion Partners: Spleen cells from immunized BALB/c mouse were fused with cells of the mouse SP2 myeloma cell line.

Specificity

This antibody recognizes rat PECAM-1 (CD31), a type 1 transmembrane protein expressed primarily on endothelial cells, platelets and leucocytes.

Formulation

Product Type: Monoclonal Antibody

Product Form: Purified IgG - liquid

Buffer Solution: Phosphate buffered saline pH7.4

Approx. Protein Concentrations: IgG concentration 0.1 mg/ml

Preservative Stabilizers: 0.09% Sodium Azide

Storage

18 months from date of dispatch.

Store at 4°C or at -20°C if preferred. This product should be stored undiluted.

Storage in frost free freezers is not recommended. Should this product contain a precipitate we recommend microcentrifugation before use.

Avoid repeated freezing and thawing as this may denature the antibody.

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**



RayBiotech, Inc.

3607 Parkway Lane suite 200
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

Applications

Table Summary of antibody applications and working conditions

| Options Functions | YES | NO | Not determined | Recommended Work dilution or concentration |
|------------------------------|------------|-----------|-----------------------|---|
| ELISA | . | | | |
| Western Blotting | | | . | |
| Immunohistology - frozen | . | | | 1/10-1/100 |
| Immunohistology - paraffin | | | . | |
| Immunohistology - resin | | | . | |
| Immunoprecipitation | | | . | |
| Flow Cytometry(*) | . | | | 1/10-1/100 |
| Immunofluorescence | | | . | |
| Neutralization | | | . | |

Note: Other applications are not tested yet. Optimal dilutions should be determined by each laboratory for each application.
(*)Use 10ul of the suggested working dilution to label 106 cells or 100ul whole blood.

Secondary Antibody Applications

| Options Functions | YES | NO | Not determined | Recommended Work dilution or concentration |
|--------------------------------------|------------|-----------|-----------------------|---|
| Immunoassay (ELISA, Western blot) | . | | | 1:5000-1:10000 |

Reference

1. Flaris, N.A. et al. (1993) Characterisation of microglia and macrophages in the central nervous system of the rat. *GLIA* 7: 34-40.
2. Male, D. et al. (1995) An interferon inducible molecule on brain endothelium which controls lymphocyte adhesion mediated by integrins. *Immunology* 84: 453-460.
3. Williams, K.C. et al. (1996) PECAM-1 (CD31) expression in the central nervous system and its role in experimental allergic encephalomyelitis in the rat. *J. Neurosci. Res.* 45(6):747-57.

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**