

RayBiotech, Inc.

3607 Parkway Lane suite 200 Norcross,GA 30092

Tel: 770-729-2992, 1-888-494-8555

Fax: 770-206-2393

Website: www.raybiotech.com Email: info@raybiotech.com

Certificate of Analysis and Data Sheet

Mouse Anti-Human Cytokeratin 19

With HRP-conjugated Secondary Antibody

Catalog No. Target Species Isotype
DS-MB-01111 Human IgG1

Preparation

Purified IgG prepared by affinity chromatography on Protein A

Immunogen: Mammary organoids

Formulation

Product Type: Monoclonal Antibody **Product Form:** Purified IgG - liquid

Buffer Solution: Phosphate buffered saline **Preservative Stabilizers:** 0.09% Sodium Azide

Approx. Protein Concentrations: IgG concentration 1.0 mg/ml

Specificity

DS-MB-01111 recognizes the 40kD cytokeratin corresponding to cytokeratin 19. Cytokeratin 19 is expressed by many simple and non-keratinizing normal and neoplastic epithelia, including glandular and ductal epithelia.

Applications

Table Summary of antibody applications and working conditions

Options Functions	YES	NO	Not determined	Recommended Work dilution or concentration
Immunohistology - Frozen	•			
Immunohistology – Paraffin (1)	•			1/50 - 1/100
Immunohistology - Resin			•	
ELISA			•	

Note: Other applications are not tested yet. Optimal dilutions should be determined. (1) This product requires protein digestion pre-treatment of paraffin sections e.g. trypsin or pronase. Histology Positive Control Tissue: Normal breast.



RayBiotech, Inc.

3607 Parkway Lane suite 200 Norcross,GA 30092

Tel: 770-729-2992, 1-888-494-8555

Fax: 770-206-2393

Website: www.raybiotech.com Email: info@raybiotech.com

Secondary Antibody Applications

Options Functions	YES	NO	Not determined	Recommended Work dilution or concentration
Immunoassay (ELISA, Western blot)	•			1:5000-1:10000

Reference

- 1. Bartek, J. et al. (1985) A subclass of luminal epithelial cells in the human mammary gland, defied by antibodies to cytokeratins. J. Cell. Sci. 75: 17-33.
- 2. Bartek, J. et al. (1985) Patterns of expression of keratin 19 as detected with monoclonal antibodies in human breast tissues and tumors. Int. J. Cancer. 36: 299-306.

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. **Shelf Life:** 18 months from date of dispatch.