



RayBiotech, Inc.

3607 Parkway Lane suite 200
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

Certificate of Analysis and Data Sheet

Mouse Anti Human HLA ABC

With HRP-conjugated Secondary Antibody

Catalog No.

DS-MB-01538

Target Species

Human

Isotype

IgG2a

Preparation

Immunogen: Purified human tonsil lymphocyte membranes.

Purification: Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.

Fusion Partners: Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1/1-Ag4.1 myeloma cell line.

Formulation

Product Type: Monoclonal Antibody

Product Form: Purified IgG - liquid

Buffer Solution: Phosphate buffered saline

Preservative Stabilizers: 0.09% Sodium Azide (NaN_3)

Approx. Protein Concentration: IgG concentration 1.0 mg/ml

Specificity

DS-MB-01538 recognizes an antigenic determinant shared among products of the HLA A, B and C loci. Clone W6/32 recognises a conformational epitope, reacting with HLA class I alpha3 and alpha2 domains ⁽⁷⁾. MHC class I glycoproteins are found on the surface of almost all nucleated somatic cells.

This product is routinely tested in flow cytometry on human peripheral blood lymphocytes.

Species Crossreactivity: Macaque, Bovine, Cat, Cynomolgus monkey and Baboon

Storage

Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody.

Shelf Life: 18 months from date of shipment

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**



RayBiotech, Inc.

3607 Parkway Lane suite 200
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

Applications

Table Summary of antibody applications and working conditions

Options Functions	YES	NO	Not determined	Recommended Work dilution or concentration
ELISA	•			
Western Blotting			•	
IHC - Paraffin		•		
IHC – Frozen (2)	•			
Immunoprecipitation	•			
Flow Cytometry (1)	•			1/50 - 1/100

Note: Other applications are not tested yet. Optimal dilutions should be determined by the user.

(1) Use 10ul of the suggested working dilution to label 10^6 cells in 100ul.

(2) Histology Positive control: Tonsil

Secondary Antibody Applications

Options Functions	YES	NO	Not determined	Recommended Work dilution or concentration
Immunoassay (ELISA, Western blot)	•			1:5000-1:10000

References

1. Barnstable, C. J. *et al.* (1978) Production of monoclonal antibodies to group A erythrocytes, HLA and other human cell surface antigens - new tools for genetic analysis. *Cell*. 14: 9–20.
2. Jacobsen, C. N. *et al.* (1993) Reactivities of 20 anti-human monoclonal antibodies with leucocytes from ten different animal species. *Vet. Immunol. Immunopathol.* 39: 461-466 .
3. Yoshino, N. *et al.* (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of Cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. *Exp. Anim.* 49: 97-110 .

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**



RayBiotech, Inc.

3607 Parkway Lane suite 200
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

4. Neefjes, J. J. *et al.* (1986) A biochemical characterization of feline MHC products: unusually high expression of class II antigens on peripheral blood lymphocytes. *Immunogenetics* 23: 341-347.
5. Stern, P. *et al.* (1987) Class I-like MHC molecules expressed by baboon placental syncytiotrophoblast. *J. Immunol.* 138: 1088-1091.
6. Verbeek, M. M. *et al.* (1995) T lymphocyte adhesion to human brain pericytes is mediated via very late antigen-4/vascular cell adhesion molecule-1 interactions. *J. Immunol.* 154: 5876-5884.
7. Tanabe, M. *et al.* (1992) Structural and functional analysis of monomorphic determinants recognized by monoclonal antibodies reacting with the HLA class I alpha 3 domain. *J. Immunol.* 148: 3203-3209.
8. Ishitani, A. *et al.* (2003) Protein expression and peptide binding suggest unique and interacting functional roles for HLA-E, F and G in maternal-placental immune recognition. *J Immunol.* 171: 1376-1384.

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**