



RayBiotech, Inc.

3607 Parkway Lane suite 200
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

Certificate of Analysis and Data Sheet

MOUSE ANTI BOVINE TNF ALPHA

With HRP-conjugated Secondary Antibody

Catalog No.
DS-MB-02436

Species
Bovine

Isotype
IgG2b

Preparation

Purification: Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Immunogen: Recombinant bovine TNF alpha

Specificity

This product recognizes bovine TNF alpha, a 17.5kDa cytokine, expressed by many different stimulated cell types including monocytes, macrophages, endothelial cells, fibroblasts and both T and B-lymphocytes.

The production of TNF alpha is induced by a variety of factors, dependant upon cell type and includes bacterial toxins, IL-1, PDGF, IFN-beta, NGF, Oncostatin M and viral infections. The presence of TNF alpha is responsible for diverse immunomodulatory, anti-tumour and toxic effects and under certain conditions is also capable of self-stimulation and inhibition.

Species Cross Reactivity: reacts with sheep (antibody reactivity and working conditions may vary between species.)

Formulation

Product Type: Monoclonal Antibody

Product Form: Purified IgG – liquid

Buffer Solution: Phosphate buffered saline pH7.4

Preservative & Stabilizers: 0.09% Sodium Azide

Approx. Protein Concentrations: IgG concentration 1.0 mg/ml

Storage

18 months from date of dispatch.

Store at 4°C or at -20°C if preferred.

This product should be stored undiluted. Storage in frost-free freezers is not recommended.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**



RayBiotech, Inc.

3607 Parkway Lane suite 200
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

Applications

Table Summary of antibody applications and working conditions

| Options Functions | YES | NO | Not determined | Recommended Work dilution or concentration |
|-----------------------------|-----|----|----------------|--|
| ELISA | • | | | |
| Western Blotting | | | • | |
| Immunohistology - frozen | | | • | |
| Immunohistology - paraffin | | | • | |
| Immunohistology - resin | | | • | |
| Immunoprecipitation | | | • | |
| Flow Cytometry(*) | • | | | |
| Immunofluorescence staining | | | • | |
| Neutralization | | | • | |

Note: Other applications are not tested yet. Optimal dilutions should be determined by each laboratory for each application.

* Use 10ul of the suggested working dilution to label 10^6 cells in 100ul

Secondary Antibody Applications

| Options Functions | YES | NO | Not determined | Recommended Work dilution or concentration |
|--------------------------------------|-----|----|----------------|--|
| Immunoassay (ELISA, Western blot) | • | | | 1:5000-1:10000 |

Reference

1. Hope, J.C. et al. (2003) Maturation of bovine dendritic cells by lipopeptides. Vet. Immunol. Immunopathol. 95: 21-31.
2. Whelan A.O. et al. (2003) Modulation of the bovine delayed-type hypersensitivity responses to defined mycobacterial antigens by a synthetic bacterial lipopeptide. Infect. Immun. 71: 6420-6425.
3. Guernon, J. et al. (2003) A tumour necrosis factor alpha autocrine loop contributes to proliferation and nuclear factor-kappaB activation of Theileria parva-transformed B cells. Cell. Microbiol. 5: 709-716.

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**