



RayBiotech, Inc.

3607 Parkway Lane suite 200
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

Certificate of Analysis and Data Sheet

Rat anti Mouse CD206:RPE

Catalog No.
DS-MB-03772

Species
Mouse

Isotype
IgG2a

Preparation

- **Preparation:** Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
- **Immunogen:** Chimaeric CRD4-7-Fc protein.
- **Fusion Partners:** Spleen cells from immunized Fischer rats were fused with cells of the Y3 myeloma cell line.

Specificity

- DS-MB-03772 recognizes the mouse mannose receptor, a 175kD type 1 membrane protein that is also known as CD206. CD206 is expressed on most tissue macrophages, *in vitro* derived dendritic cells, lymphatic and sinusoidal endothelia.
- DS-MB-03772 has been reported to be non-inhibitory for the binding of the mannose receptor to carbohydrate ligands.

Formulation

- **Product Type:** Monoclonal Antibody
- **Product Form:** Purified IgG conjugated to R. Phycoerythrin (RPE) – lyophilized
- **Reconstitution:** Reconstitute in 0.25 ml distilled water
- **Buffer Solution:** Phosphate buffered saline, pH 7.4
- **Preservative & Stabilizers:** 0.09% Sodium Azide; 1% Bovine Serum Albumin
- **Approx. Protein Concentrations:** IgG concentration 0.1 mg/ml

Storage

- Store at 4°C. **DO NOT FREEZE**
- This product should be stored undiluted. This product is photosensitive and should be protected from light.
- **Shelf Life:** 12 months from date of shipment.

Reference:

- Martinez-Pomares, L. et al. (2003) Analysis of mannose receptor regulation by IL-4, IL-10 and proteolytic processing using novel monoclonal antibodies. J. Leuk. Biol. 73: 604 - 613.

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**



RayBiotech, Inc.

3607 Parkway Lane suite 200
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

- Zamze, S. et al. (2002) Recognition of bacterial capsular polysaccharides and lipopolysaccharides by the macrophage mannose receptor. J. Biol. Chem. 277: 41613 - 41623.

Applications

Options Functions	YES	NO	Not determined	Recommended Work dilution or concentration
ELISA			•	
Flow Cytometry(1)	•			Neat

Note: Other applications are not tested yet. Optimal dilutions should be determined by each laboratory for each application.
(1) Use 10ul of the suggested working dilution to label 10^6 cells in 100ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**