

### RayBiotech, Inc.

3607 Parkway Lane suite 200 Norcross,GA 30092

Tel: 770-729-2992, 1-888-494-8555

Fax: 770-206-2393

Website: www.raybiotech.com Email: info@raybiotech.com

# Certificate of Analysis and Data Sheet

# Rabbit Anti-Human Collagen III

With HRP-conjugated Secondary Antibody

Catalog No. Target Species Isotype
DS-PB-00518 Human Polyclonal IgG

## Preparation

Immunogen: Collagen III from human placenta

**Antiserum Preparation:** Antiserum to native and denatured human collagen type III was raised by repeated immunization of rabbits with highly purified antigen. DS-PB-00518 has been cross-absorbed

with immobilized human collagen types I, IV and V and with human blood plasma proteins.

#### **Formulation**

**Product Type:** Polyclonal Antibody **Product Form:** Purified IgG - lyophilized

**Reconstitution:** Reconstitute with 0.5 ml distilled water

\* For long term storage the addition of 0.09% sodium azide is recommended.

**Buffer Solution:** Mannitol, salts and dextran **Preservative Stabilizers:** See buffer solution

**Approx. Protein Concentrations:** 0.1 mg/ml after reconstitution

## Specificity

DS-PB-00518 is specific for human collagen type III. It has shown the following cross-reactivities.

Human collagen type I <15% Human collagen type IV <5% Human collagen type II and V <1%

Human plasma proteins do not interfere with binding of the antibody to collagen.

### Storage

Prior to reconstitution store at +4°C.

After reconstitution store at -20°C.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody.

**Shelf Life:** 12 months from date of reconstitution.

#### References

1. George, C. L. *et al* (2001) Endotoxin responsiveness and subchronic grain dust-induced airway disease. Am. J. Physiol. Lung Cell. Mol. Physiol. 280(2): 203-213.



## RayBiotech, Inc.

3607 Parkway Lane suite 200 Norcross,GA 30092

Tel: 770-729-2992, 1-888-494-8555

Fax: 770-206-2393

Website: www.raybiotech.com Email: info@raybiotech.com

**Applications** 

Options Functions	YES	NO	Not determined	Recommended Work dilution or concentration
Immunohistology - Frozen	•			
Immunohistology - Paraffin	•			1/10 - 1/40
Immunohistology - Resin			•	
ELISA	•			1/2000 - 1/8000
Western Blotting			•	

Note: Other applications are not tested yet. Optimal dilutions should be determined.

## Secondary Antibody Applications

Options Functions	YES	NO	Not determined	Recommended Work dilution or concentration
Immunoassay (ELISA, Western blot)	•			1:5,000-1:10,000

## References

1. George, C. L. *et al* (2001) Endotoxin responsiveness and subchronic grain dust-induced airway disease. Am. J. Physiol. Lung Cell. Mol. Physiol. 280(2): 203-213.