Bsp19 I

C^CATGG AccuCut™ Restriction Endonuclease

• Cat. No. E-1381 500 Units E- 1382 2500 Units

Lot No.: 02I131491H8A3

· Supplied with Enzyme

10X	AccuCut™B lu	e Buffer :	1 mL
	100 mM	pH 8.5	Tris-HC
	100 mM	MgCl ₂	
	1 M	NaCl	
	10 mM	DTT	
1X Di	lution Buffer	:	1 mL
	10 mM	pH 7.6	Tris-HC
	50 mM	KCI	
	0.1 mM	EDTA	
	1 mM	DTT	
	200μ g/mL	Acetyl	ated BSA
	50%	Glycer	ol

- Store at -20 C.
- Unit definition : One unit of restriction endonuclease activity is defined as the amount of enzyme required to completely digest 1µg of substrate DNA in a total reaction volume of 50 µL in one hour using the AccuCut™ buffer provided. Incubations are performed in 1.5 mL tubes at the appropriate incubation temperature as indicated in the Product Profile.

• Isoschizomer : Nco I.

• Neoschizomer: Unfound

 Reactivity on methylated substrate DNA: unidentified

• Ref) 1. Degtyarev, S.K., Unpublished observations

Source : Bacillus species 19.

· Concentration: 20 Units/uL

Reaction Condition

- 10X AccuCut™ Blue Buffer

- Incubate at 37℃.

Storage Buffer

 20 mM
 pH 7.5, Tris-HCI

 50 mM
 KCI

 1 mM
 EDTA

 10 mM
 2-mercaptoethanol

 50%
 Glycerol

Heat inactivation: 65 °C for 20 minutes.

Quality Control

· Overdigestion Assay :

No nonspecific activity was detected after incubation of 1 μg of λ DNA with 50 units of Bsp19 I for 15 hours.

- * Conditions of low ionic strength, high enzyme concentration, glycerol concentration >5%, or pH >8.0 may result in star activity.
- Nuclease Contamination Assay :

No altered pattern was detected after incubation of 1 μg of substrate DNA with *Bsp*19 I in 50 μL reaction volume with the supplied AccuCut™ buffer overnight.

· Ligation and Recutting Assay:

This assay is used to test for exonuclease activity that would degrade the termini of restriction fragments, resulting in inhibition of ligation and of subsequent digestion of ligated fragments. After 40-fold overdigestion with *Bsp*19 I, 95% of the DNA fragments can be ligated and recut with *Bsp*19 I.