

Olig2

Data Sheet

| | | | |
|----------------------------|--|----------------------------|---|
| Catalog Number: | GT15132 | Host: | Goat |
| Product Type: | Affinity purified | Species Reactivity: | Human |
| Immunogen Sequence: | <i>E. coli</i> -derived, recombinant human Oligodendrocyte transcription factor 2 (rhOlig2). | Format: | Liquid 1mg/ml Solution in phosphate-buffered saline (PBS) with 5% Trehlose |
| Applications: | Immunohistochemistry : 10 µg/mL Western Blot: 0.1 - 0.2 µg/mL ELISA: 0.5 - 1.0 µg/mL | | |

Dilutions listed as a recommendation. Optimal dilution should be determined by investigator.

Storage: Antibody can be aliquotted and stored frozen at -20° C to -70° C in a manual defrost freezer for six months without detectable loss of activity. The antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. *Avoid repeated freeze-thaw cycles.*

Application Notes

Direct ELISA

This antibody can be used at 0.5 - 1.0 µg/mL with the appropriate secondary reagents to detect human Olig2. The detection limit for rhOlig2 is approximately 0.3 ng/well. In this format, this antibody shows approximately 10% cross-reactivity with rhOlig3 and 5% cross-reactivity with rhOlig1.

Western blot

This antibody can be used at 0.1 - 0.2 µg/mL with the appropriate secondary reagents to detect human Olig2. The detection limit for rhOlig2 is approximately 5 ng/lane and 1 ng/lane under non-reducing and reducing conditions, respectively.

Immunocytochemistry

This antibody has been used at a concentration of 10 µg/mL to detect Olig2 on mouse E11 spinal cord tissue sections. Sections were fixed with PBS containing 4% paraformaldehyde for 20 minutes at room temperature and blocked with PBS containing 10% normal donkey serum, 0.1% Triton X-100 and 1% BSA for 45 minutes at room temperature. After blocking, cells were incubated with diluted primary antibody overnight at 4° C followed by Rhodamine Red coupled anti-goat IgG at room temperature in the dark for one hour. Between each step, cells were washed with PBS containing 0.1% BSA.

FOR RESEARCH USE ONLY

NEUROMICS' REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE.

6/04v1