



Hepatitis B Virus Protein X Hepatitis B virus E. coli Tag free

Product Data Sheet

Type: Recombinant

Tag: Tagless

Source: E. coli

Species: Hepatitis B virus

Other names: HBx

Cat. No.:

RD972038100 (0.1 mg)

Description

Total 154 AA. MW: 17 kDa (calculated). 153 AA of recombinant HBx protein and one extra AA, N-terminal methionin (highlighted).

Introduction to the Molecule

Hepatitis B virus X protein (HBx) is a 17 kD transcriptional coactivator that plays a major role in the regulation of genes related to inflammation and cell survival. It regulates many transcription factors including nuclear factor kappa B (NF-kappaB). It also plays a key role in hepatocarcinogenesis. HBx facilitates the binding of cAMP response element binding protein (CREB) to its responsive element. HBx stabilizes the cellular coactivator ASC-2 through direct protein-protein interaction, affecting the regulation of genes, which are actively transcribed in liver cancer cells.

Moreover, HBx transactivates both JNK and MAPK signal transduction pathways in association with the mobilization of cytosolic Ca²⁺. In addition, HBx decreases the expression of PTEN a known tumor suppressor and a negative regulator of phosphatidylinositol 3'-kinase/AKT. HBx also decreased the expression of PTEN in HBx-transfected cells.

The etiology of hepatocellular carcinoma (HCC) is involved with hepatitis B virus (HBV) infection and HBx in particular plays a role in the development of HBV-related HCC. The persistence of HBx is important to the pathogenesis of early HCC. HBx expression in the liver during chronic HBV infection may be an important prognostic marker for the development of HCC.

Research topic

Others

Amino Acid Sequence

MAARVCCQLD PARDVLCLRP VGAESRGRPV SGPFGTLPSP SSSAVPADHG AHLRLRGLPV CAFSSAGPCA LRFTSARRME
TTVNAHQVLP KVLHKRTLGL SAMSTTDLEA YFKDCLFKDW EELGEEIRLK VFVLGGCRHK LVCSPAPCNF FTSA


Source

E. coli

Purity

>98%

SDS-PAGE gel



12% SDS-PAGE separation of HBx

1. M.W. marker - 14, 21, 31, 45, 66, 97 kDa

2. reduced and heated sample, 5µg/lane

3. non-reduced and non-heated sample, 5µg/lane

Endotoxin

<1.0 EU/µg

Formulation

Filtered (0.4 µm) and lyophilized in 0.5 mg/mL in 0.05 M phosphate buffer, 0.075 M NaCl, pH 7.4

Reconstitution

Add deionized water to prepare a working stock solution of 0.5 mg/mL and let the lyophilized pellet dissolve completely.

Product is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.

Shipping

At ambient temperature. Upon receipt, store the product at the temperature recommended below.

Storage, Stability/Shelf Life

Store lyophilized protein at -80°C. Lyophilized protein remains stable until the expiry date when stored at -80°C. Aliquot reconstituted protein to avoid repeated freezing/thawing cycles and store at -80°C for long term storage. Reconstituted protein can be stored at 4°C for a week.

Quality Control Test

BCA to determine quantity of the protein.

SDS PAGE to determine purity of the protein.

LAL TEST to determine quantity of endotoxin.

Applications

ELISA, Western blotting

Note

This product is intended for research use only.

Gentaur Molecular Products
Voortstraat 49
1910 Kampenhout, Belgium
<http://www.gentaur-worldwide.com>