DATA SHEET

Fluorescein (FITC) conjugated Affinity pure Goat anti-Human IgG (H+L) (Minimum cross reactivity with bovine, horse, mouse, and rabbit serum proteins)

Catalog number: SA-9001-15

Description: This affinity pure anti-human immunoglobulin was purified from healthy goat immunized with purified human IgG (H+L). The immunized serum was delipidated and heat inactivated at 56°C before affinity purification. The purified IgG (H+L) was treated with pH 2.8 for at least 30 minutes.

Size: 1 mg/1 ml

Antibody Concentration: One mg of affinity pure antibody was conjugated with Fluoroscein isothiocyanate (FITC); the conjugate was purified on gel filtration column.

 $\textbf{Fluorophore:} \ \mathsf{Fluoroescein-5-isothiocyanate} \ (\mathsf{FITC} \ isomer 1)$

Absorbance max. = 495 nm; E max. = 520 nm

A 495 nm / A280 nm = 1.1

.Intended Use: Immunofluorescence 1:50-1:200

Flow cytometry 1:1,000 1:10,000 (not tested in our lab).

The optimum dilution should be determined by the individual lab.

Storage: Storage: 2-8°C

Buffer: 10 mM phosphate, 150 mM NaCl, pH 7.6, 1 % BSA (IgG and protease free) and 0.05% sodium

azide

Antibody specificity: The antibody is purified on immunoaffinity human IgG (H+L) column. Using ELISA, this antibody reacts with human IgG (H+L), heavy and light chains and light chains of other human immunoglobulins. This antibody does not cross react with non-immunoglobulin serum proteins. This antibody has minimum cross reactivity with bovine, horse, mouse and rabbit serum proteins. Cross reactivity to other serum proteins have not been determined.

Limitation and warranty: Our warranty is limited to the actual price paid for the product. We are not liable for any property damage, personnel injury, time, effort or economic loss due to our product.

MSDS: This product contains 0.05 % sodium azide as a preservative, appropriate care should be taken in handling. National Institute of Occupational Safety and Health has warning that sodium azide can react with lead, copper, brass or solder in the plumbing system and forms hydrazoic acid in acidic condition. Discard with copious amount of water. Avoid skin and eye contact with all laboratory products. Use appropriate lab. gear, lab coat, gloves and safety glasses. Do not ingest any lab. products. This product is not approved for administration in human or animals.

For research use only; not for use in diagnostic procedures. FOR IN VITRO LABORATORY USE ONLY

PRODUCT OF USA