

bs-0552R-Cy7

• Rabbit Anti-Collagen V Polyclonal Antibody, Cy7 conjugated

Conjugated Primary Antibodies

Background:

Collagens are highly conserved throughout evolution and are characterized by an uninterrupted "Glycine-X-Y" triplet repeat that is a necessary part of the triple helical structure. Type V Collagen is a member of group I Collagen (fibrillar forming collagen). It is a minor connective tissue component of nearly ubiquitous distribution. Type V Collagen binds to DNA, heparan sulfate, Thrombospondin, Heparin, and Insulin. Collagen V is composed of several subunits as follows: trimers of two alpha 1(V) and one alpha 2(V) chains in most tissues and trimers of one alpha 1(V), one alpha 2(V), and one alpha 3(V) chains in placenta. Type V Collagen is closely related to type XI Collagen and it is possible that the Collagen chains of types V and XI constitute a single collagen type with tissue specific chain combinations. Mutations in this gene are associated with Ehlers Danlos syndrome, types I and II.

Purification: Was purified by Protein A and peptide affinity chromatography.



Storage:

Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year. Protect from light.

Reconstitution:

If the antibody is in liquid form, it is ready to use, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.



For full size images and description please click [HERE](#).

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities:

Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Rabbit, Sheep,

Application:

- IF (1:100-500)
- Not yet tested in other applications. Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 202kDa

Preservatives: 10ug/uL BSA and 0.1% NaN₃.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.