

bs-0552R-FITC

• Rabbit Anti-Collagen V Polyclonal Antibody, FITC conjugated

Conjugated Primary Antibodies

Background:

Collagens are highly conserved throughout evolution and are characterized by an uninterrupted "Glycine-X-Y" triplet repeat that is a necessary part of the triple helical structure. Type V Collagen is a member of group I Collagen (fibrillar forming collagen). It is a minor connective tissue component of nearly ubiquitous distribution. Type V Collagen binds to DNA, heparan sulfate, Thrombospondin, Heparin, and Insulin. Collagen V is composed of several subunits as follows: trimers of two alpha 1(V) and one alpha 2(V) chains in most tissues and trimers of one alpha 1(V), one alpha 2(V), and one alpha 3(V) chains in placenta. Type V Collagen is closely related to type XI Collagen and it is possible that the Collagen chains of types V and XI constitute a single collagen type with tissue specific chain combinations. Mutations in this gene are associated with Ehlers Danlos syndrome, types I and II.

Purification: Was purified by Protein A and peptide affinity chromatography.

Storage:

Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year. Protect from light.

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities:

Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Rabbit, Sheep,

Application:

- IF (1:100-500)
- Not yet tested in other applications. Optimal working dilutions must be determined by the end user.



Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 202kDa

Preservatives: 10ug/uL BSA and 0.1% NaN₃.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.



For full size images and description please click [HERE](#).