www.biossusa.com support@biossusa.com 800.501.7654 [DOMESTIC] +1.781.569.5821 [INTERNATIONAL]

# Bioss

# bs-0641R-Cy7

# Rabbit Anti-Integrin alpha 4/CD49d/Integrin a4 Polyclonal Antibody, Cy7 conjugated

Conjugated Primary Antibodies

#### Background:

Integrin alpha 4 (also called CD49d) is a 150 kDa protein that possesses a large extracellular domain involved in ligand binding, a single transmembrane domain, and an intracellular regulatory domain possessing multiple sites for phosphorylation. Integrin alpha 4 forms heterodimers with integrins beta 1 and beta 7. Integrin alpha 4 is expressed on leukocytes and leukocyte precursors, neural crest cells, and developing skeletal muscles and is essential for embryogenesis, hematopoiesis, and immune responses. The presence of integrin alpha 4 promotes cell migration and inhibits cell spreading and contractility. Integrin alpha 4 function has been implicated in the pathogenesis of multiple diseases including asthma, rheumatoid arthritis, Crohn's disease, ulcerative colitis, hepatitis C, and multiple sclerosis, and therefore, modulation of integrin alpha 4 function has become an important target for drug discovery.

Purification: Was purified by Protein A and peptide affinity chromatography.

#### Storage

Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year. Protect from light.

### Reconstitution:

If the antibody is in liquid form, it is ready to use, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

For full size images and description please click  $\ensuremath{\mathsf{HERE}}\,.$ 

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit Reactivities:

Human, Mouse, Rat, Cow, Horse, Rabbit,

## Application:

FACS(1:100-500)IF(1:100-500)

 Not yet tested in other applications.
Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 114kDa

Preservatives:

10ug/uL BSA and 0.1% NaN3.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.