bs-0777R

· Rabbit Anti-Mink IgG Antibody (H+L). Unconjugated

Secondary Antibodies

Background:

Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 1011 variants.

Purification: Was purified by Protein A and peptide affinity chromatography.

Storage

Prepared as lyophilized powder and shipped on ice. Store at -20°C for one year as lyophilized powder or liquid. Please reconstitute before use.

Reconstitution:

Two additional vials are included in shipment for reconstitution purposes (double distilled H20 and sterile glycerol). Centrifuge all vials to ensure necessary quantities have settled. Add 50uL of sterile double distilled water to antibody. Mix thoroughly by gently pipetting up and down. Then, add 50uL of sterile glycerol to antibody. Mix well and keep cold. *Please note lyophilized powder is supplied in bovine serum albumin and sodium azide. After reconstitution, final concentration of preservatives are 10ug/uL BSA and 0.1% NaN3.

Gentaur Molecular Products Voortstraat 49 1910 Kampenhout, Belgium Size: 1mg lyophilized

Concentration: 10ug/1uL

Application:

- WB
- ELISA
- IP
- IHC • ICC
- Not yet tested in other applications.
 Optimal working dilutions must be determined by the end user.

Cross-reactivities: Mink

Molecular Weight: 150kDa

Isotype: IgG (H+L)

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.