### bs-1301R

# Rabbit Anti-FGFR3/CD333 Polyclonal Antibody

spliced transcript variants encoding the same protein have been described.

Primary Antibodies

#### Background:

The encoded protein is synthesized mainly in corticotroph cells of the anterior pituitary where four deavage sites are used; adrenocorticotrophin, essential for normal steroidogenesis and the maintenance of normal adrenal weight, and lipotropin beta are the major end products. In other tissues, including the hypothalamus, placenta, and epithelium, all cleavage sites may be used, giving rise to peptides with roles in pain and energy homeostasis, melanocyte stimulation, and immune modulation. These include several distinct melanotropins, lipotropins, and endorphins that are contained within the adrenocorticotrophin and beta-lipotropin peptides. Mutations in this gene

#### Source/Purification:

KLH conjugated synthetic peptide derived from human FGFR3. Was purified by Protein A and peptide affinity chromatography.

have been associated with early onset obesity, adrenal insufficiency, and red hair pigmentation. Alternatively

Storage: Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year.

## Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities:

Human, Mouse, Rat, Chicken, Dog, Cow, Horse, Sheep,

## Application:

- ELISA(1:500-1000)
- IP(1:20-100)
- IHC-P(1:100-500)
- IHC-F(1:100-500)
  IF(1:100-500)
- Not yet tested in other applications.
  Optimal working dilutions must be determined by the end user.

Antibody Type: Polydonal

Isotype: IgG

Molecular Weight: 86kDa

Preservatives:

10ug/uL BSA and 0.1% NaN3.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.