Rabbit Anti-Phospho-Smad1(Ser206) Polyclonal Antibody

Primary Antibodies

Background:

The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signals of the bone morphogenetic proteins (BMPs), which are involved in a range of biological activities including cell growth, apoptosis, morphogenesis, development and immune responses. In response to BMP ligands, this protein can be phosphorylated and activated by the BMP receptor kinase. The phosphorylated form of this protein forms a complex with SMAD4, which is important for its function in the transcription regulation. This protein is a target for SMAD-specific E3 ubiquitin ligases, such as SMURF1 and SMURF2, and undergoes ubiquitination and proteasome-mediated degradation. Alternatively spliced transcript variants encoding the same protein have been observed. [provided by RefSeq].

Source/Purification:

KLH conjugated Synthesised phosphopeptide derived from human Smad1 around the phosphorylation site of Ser206. Was purified by Protein A and peptide affinity chromatography.

Modification Site:

Ser206

Storage: Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year.

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities: Human, Mouse, Rat,

Application:

WB(1:100-500)

ELISA(1:500-1000)

IP(1:20-100)

IHC-P(1:100-500)

IHC-F(1:100-500)

• IF(1:100-500)

 Not yet tested in other applications.
Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 52kDa

Preservatives:

10ug/uL BSA and 0.1% NaN3.

For research use only, CAUTION: Not for human or animal therapeutic or diagnostic use.