

bs-7694R-HRP

• Rabbit Anti-SLC12A3/NCCT Polyclonal Antibody, HRP conjugated

Conjugated Primary Antibodies

Background:

Na-K-Cl cotransporters (NKCC) are channel proteins that aid in the transcellular movement of chloride across both secretory and absorptive epithelia. NKCC1 is expressed in muscle cells, neurons, and red blood cells. In the basolateral membrane of secretory epithelia, NKCC1 mediates active chloride secretion. The gene encoding human NKCC1 maps to chromosome 5q23.3. In mice, disruption of the NKCC1 gene leads to deafness and impaired balance. NKCC2 is specifically expressed in the kidney where it mediates active reabsorption of sodium chloride in the thick ascending limb of the loop of Henle. NKCC2 is sensitive to the clinically important diuretics furosemide and bumetanide. The gene encoding human NKCC2 maps to chromosome 15q15-q21 and mutations in this gene lead to Bartter's syndrome, an inherited hypokalaemic alkalosis. NCCT is a thiazide-sensitive Na-Cl cotransporter that is primarily expressed in the distal convoluted tubule of the kidney where it accounts for a significant fraction of net renal sodium reabsorption. The gene for human NCCT map to chromosome 16q13. Mutations in the gene encoding NCCT cause Gitelman's syndrome, a subset of Bartter's syndrome.

Purification: Was purified by Protein A and peptide affinity chromatography.

Storage: Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year.

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities:

Human, Mouse, Rat, Dog, Cow, Rabbit,

Application:

- WB(1:100-500)
- ELISA(1:500-1000)
- IHC-P(1:100-500)
- IHC-F(1:100-500)
- Not yet tested in other applications. Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 113kDa

Preservatives:

10ug/uL BSA and 0.01% Gentamicin.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.

For full size images and description please click [HERE](#).