

bs-9856R-FITC

• Rabbit Anti-HCN3 Polyclonal Antibody, FITC conjugated

Conjugated Primary Antibodies

Background:

Hyperpolarization-activated, cyclic nucleotide-binding channels (HCN) are voltage-gated cation channels that are activated by direct binding of intracellular cyclic nucleotides. The HCN family consists of four members (HCN1-4), each with a core transmembrane segment domain and a C-terminal 120 amino-acid cyclic nucleotide-binding domain motif. HCN channels are expressed in the brain, heart, thalamus and testis. The pacemaker properties of HCN channels contribute to spontaneous rhythmic activity in the brain and heart. HCN3 contains a segment characterized by a series of positively charged amino acids at every third position. This region designated S4 is likely to be the voltage sensor of the protein. In the brain, HCN3 and HCN4 exhibit subcortical distribution mainly concentrated in the hypothalamus and thalamus, respectively.

Purification: Was purified by Protein A and peptide affinity chromatography.

Storage:

Aqueous buffered solution containing 100ug/ml BSA, 50% glycerol and less than 0.09% sodium azide. Store at -20°C for 12 months. Protect from light. [Product without BSA and/or sodium azide is available for special order.]

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

Size: 100ul

Concentration: 1ug/uL

Host: Rabbit

Reactivities:

Human, Mouse, Rat, Bovine, Rabbit,

Application:

- IF(1:50-200)
- Not yet tested in other applications. Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 86kDa

Note:

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.

For full size images and description please click [HERE](#).