www.biossusa.com support@biossusa.com 800.501.7654 [DOMESTIC] +1.781.569.5821 [INTERNATIONAL]



bs-9907R-FITC

Rabbit Anti-CPN2 Polyclonal Antibody, FITC conjugated

Conjugated Primary Antibodies

Background:

CPN2 is a zinc metalloprotease, and cleaves carboxy-terminal arginines and lysines from peptides found in the bloodstream such as complement anaphylatoxins, kinins, and creatine kinase MM (CK-MM). By removing only one amino acid, CPN has the ability to change peptide activity and receptor binding. It is a 280 kDa tetrameric glycoprotein that is synthesized by the liver and secreted into the plasma. It consists of 2 identical 83 kDa regulatory subunits (CPN2) and 2 identical 50 kDa catalytic subunits (CPN1). CPN2, the 83 kDa subunit, binds and stabilizes the catalytic subunit at 37 degrees Celsius and keeps it in circulation. Under some circumstances it may be an allosteric modifier of the catalytic subunit. CPN is a member of a larger family of carboxypeptidases, many of which also cleave arginine and lysine. Because of the highly conserved active sites and the possible redundant functions of carboxypeptidases, it has been difficult to elucidate the role of CPN in disease processes.

Purification: Was purified by Protein A and peptide affinity chromatography.

Storage:

Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year. Protect from light.

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

For full size images and description please click HERE.

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit **Reactivities:**

Human, Mouse, Rat, Dog, Pig, Cow, Horse, Rabbit, Sheep,

Application:

- IF(1:50-200)
- Not yet tested in other applications.
 Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 58kDa

Preservatives: 10ug/uL BSA and 0.1% NaN3.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.