www.biossusa.com support@biossusa.com 800.501.7654 [DOMESTIC] +1.781.569.5821 [INTERNATIONAL]

Bioss

bsm-0917M-Cy5

Mouse Anti-BrdU(A7) Monoclonal Antibody, Cy5 conjugated

Conjugated Primary Antibodies

Background:

Bromodeoxyuridine (BrdU) is a thymidine analog and is specifically incor-porated into DNA during DNA synthesis. Anti-bromodeoxyuridine monoclonal antibody is used to identify cells that have incorporated BrdU. This immunological detection scheme has several advantages over the use of radioactive thymidine incorporation for identifying cells under-going replication. Labeling and detection can be performed the same day instead of waiting several days, as required for autoradiography of tritium-labeled cells, and the necessity of using multiple specimens for obtaining the optimal exposure time is eliminated. In addition, anti-bromodeoxyuridine staining with flow cytometric analysis allows multiple parameters to be evaluated simultaneously. Anti-bromodeoxyuridine monoclonal antibody has been used for identi-fying proliferating cells in blood (Campana et al., 1988), tissues (Schutte et al., 1987; Hayashi, et al., 1988), tumors (Hoshino et al., 1986; Morstyn et al., 1983), as well as for determining plasma cell labeling indices (Greipp et al., 1985).

Purification: Was purified by Protein A and peptide affinity chromatography.


Storage:


Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year. Protect from light.

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.





For full size images and description please click HERE.

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Mouse

Reactivities: BrdU

Application:

• IF(1:100-500)

 Not yet tested in other applications.
Optimal working dilutions must be determined by the end user.

Antibody Type: Monoclonal

Isotype: IgG

Molecular Weight: 0.3071kDa

Preservatives:

10ug/uL BSA and 0.1% NaN3.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.