



11 Park Drive, Suite 12
Boston, MA 02215

Rat monoclonal anti mouse CD11b (Azide-free & Low endotoxin)

ORDERING INFORMATION

Catalog Number:	gAP-0006
Size:	1.00 mg
Storage:	< -20° C
Immunogen:	Mouse CD11a⁺ T Cells
Ig Type:	Rat IgG2b
Clone	AP-MAB0813
Endotoxin Level	< 0.002EU/μg IgG*
Applications:	Blocking of CD11b mediated cell adhesion

Description: CD11b is a 170 kD glycoprotein, also known as α_M integrin, Mac-1 α subunit, Mol, CR3, or Ly-40. CD11b is a member of the integrin family, primarily expressed on granulocytes, monocytes/macrophages, NK cells, and some T and B cells. CD11b non-covalently associates with CD18 (β_2 integrin) to form Mac-1. Mac-1 plays an important role in cell-cell interaction by binding its ligands ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4 (CD242), iC3b, and fibrinogen.

Preparation: This antibody was produced from a hybridoma (mouse myeloma fused with spleen cells from a Rat immunized with mouse CD11b⁺ T cells).

Formulation: The IgG fraction of **culture supernatant** was purified by Protein A/G affinity chromatography and lyophilized from a 0.2 μm filtered solution in phosphate-buffered saline (PBS, **Azide Free**).

Reconstitution: Reconstitute the antibody with sterile PBS and the reconstituted antibody can be aliquoted and stored frozen at < -20 for at least for six months without detectable loss of activity. **Avoid repeated freeze-thaw cycles.** Lyophilized samples are stable for 2 years from date of receipt when stored at -70°C.

Specificity: The antibody was selected for its ability to detect mouse CD11b⁺ T cells by FC.

***Endotoxin Level:** Extremely low level of LPS (< 0.002EU/μg IgG)

Application(s):

- | | | |
|----|--|------------|
| 1. | Blocking of CD11b mediated cell-cell adhesion | Yes |
|----|--|------------|

*** The antibody is produced by in vitro culture.**

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Tel: 617-549-2665; Fax: (480) 247-4337, angioproteomie@gmail.com