




11 Park Drive, Suite 12
Boston, MA 02215

Rabbit anti Bcr (pTyr177)

- Catalog Number:** pAP11199
Amount: 100µl (for 10 Western Blotting)
Swiss-Prot No. : P11274
Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage/Stability: Store at -20°C/1 year
Immunogen: The antiserum was produced against synthesized phosphopeptide derived from human Bcr around the phosphorylation site of tyrosine 177 (P-F-Y^P-V-N).
Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.
Specificity/Sensitivity: Bcr (phospho-Tyr177) antibody detects endogenous levels of Bcr only when phosphorylated at tyrosine 177.
Reactivity: Human, Mouse
Applications: WB: 1:1000
References: Sattler M, et al. Mol Cell Biol. 1999 Nov; 19(11): 7473-7480.
Li S et al. EMBO J. 2001 Dec 3; 20(23): 6793-6804.
Million RP, et.al. Mol Cell Biol. 2004 Jun; 24(11): 4685-4695.
Malcolm A. Meyn, et al. . Biol. Chem., Oct 2006; 281: 30907 – 30916.



Western blot analysis of extract from A431 cells, untreated or treated with EGF (200ng/ml, 5min), using Bcr (Ab-177) antibody (AP21197, Lane 1 and 2) and Bcr(phospho-Tyr177) antibody (pAP11199, Lane 3 and 4).

For Research Use Only

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Fax: (480) 247-4337, angioproteomie@gmail.com