



11 Park Drive, Suite 12  
Boston, MA 02215

## Human Fibroblast Growth Factor-Acidic (FGF-1, Sf9)

### ORDERING INFORMATION

Catalog No: rAP-0018;

Size: 2 µg; 10 µg

Storage: <- 20° C

### Synonyms:

HBGF-1, ECGF-beta, FIBP, FGFIBP, FIBP-1, ECGF, ECGFA, GLIO703, FGF1, FGF-a.

### Introduction:

Acidic fibroblast growth factor is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein functions as a modifier of endothelial cell migration and proliferation, as well as an angiogenic factor. It acts as a mitogen for a variety of mesoderm- and neuroectoderm-derived cells in vitro, thus is thought to be involved in organogenesis. Three alternatively spliced variants encoding different isoforms have been described. The heparin-binding growth factors are angiogenic agents in vivo and are potent mitogens for a variety of cell types in vitro. There are differences in the tissue distribution and concentration of these 2 growth factors.

### Description:

Fibroblast Growth Factor-1 Human Recombinant (FGF-1) produced in Sf9 insect cells is a single, glycosylated, polypeptide chain containing 140 amino acids and having a molecular mass of 15803 Dalton. The FGF-a is purified by proprietary chromatographic techniques.

### Source:

*Baculovirus*

### Physical Appearance:

Sterile Filtered liquid formulation.

### Formulation:

The sterile protein solution (1.8mg/ml) contains 20mM Tris HCl pH=7.9, 100mM KCl, 0.2mM DTT and 20% glycerol.

### Stability:

Fibroblast Growth Factor-acidic although stable at 4°C for 3 weeks, should be stored desiccated below -18°C.

For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

**Please prevent freeze-thaw cycles.**

### Purity:

Greater than 95.0% as determined by:

(a) Analysis by RP-HPLC.

(b) Analysis by SDS-PAGE.

### Amino acid sequence:

The sequence of the first five N-terminal amino acids as determined and was found to be Met-Phe-Asn-Leu-Pro.

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Fax: (480) 247-4337, [angioproteomie@gmail.com](mailto:angioproteomie@gmail.com)



11 Park Drive, Suite 12  
Boston, MA 02215

**Biological Activity:**

The ED50, calculated by the dose-dependant proliferation of BAF3 cells expressing FGF receptors (measured by  $^3\text{H}$ -thymidine uptake) is <10 ng/ml, corresponding to a specific activity of  $10^5$  Units/mg.

**Usage:**

Angio-Proteomie's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Fax: (480) 247-4337, [angioproteomie@gmail.com](mailto:angioproteomie@gmail.com)