



11 Park Drive, Suite 12
Boston, MA 02215

Mouse Fibroblast Growth Factor-Acidic (FGF-1)

ORDERING INFORMATION

Catalog No: rAP-0019;

Size: 10 µg; 50 µg

Storage: <- 20° C

Synonyms:

HBGF-1, ECGF-beta, FIBP, FGFIBP, FIBP-1, ECGF, ECGFA, GLIO703, FGF1, FGF-a.

Introduction:

Acidic fibroblast growth factor is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein functions as a modifier of endothelial cell migration and proliferation, as well as an angiogenic factor. It acts as a mitogen for a variety of mesoderm- and neuroectoderm-derived cells in vitro, thus is thought to be involved in organogenesis. Three alternatively spliced variants encoding different isoforms have been described. The heparin-binding growth factors are angiogenic agents in vivo and are potent mitogens for a variety of cell types in vitro. There are differences in the tissue distribution and concentration of these 2 growth factors.

Description:

Fibroblast Growth Factor-acidic Mouse Recombinant (FGF-1) produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 141 amino acids and having a molecular mass of 15.9 kDa. The FGF acidic is purified by proprietary chromatographic techniques.

Source:

Escherichia Coli.

Physical Appearance:

Sterile Filtered White Lyophilized (freeze-dried) powder.

Formulation:

Lyophilized at a concentration of 1 mg/ml in 5mM NaP, pH-7.5, 50mM NaCl.

Solubility:

It is recommended to reconstitute the lyophilized Fibroblast Growth Factor-acidic in sterile 18MQ-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Stability:

Lyophilized Fibroblast Growth Factor-1 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution FGF-a should be stored at 4°C between 2-7 days and for future use below -18°C.

Please prevent freeze-thaw cycles.

Purity:

Greater than 98.0% as determined by:

(a) Analysis by RP-HPLC.

(b) Analysis by SDS-PAGE.

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Fax: (480) 247-4337, angioproteomie@gmail.com



11 Park Drive, Suite 12
Boston, MA 02215

Amino acid sequence:

MFNLPLGNYK KPKLLYCSNG GHFLRILPDG TVDGTDRSD QHIQLQLSAE SAGEVYIKGT
ETGQYLAMDT EGLLYGSQTP NEECLFLERL EENHYNTYTS KKHAENWFV GLKKN GSCKR
GPRTHYGQKA ILFLPLPVSS D.

Biological Activity:

The ED₅₀ as determined by the dose-dependent stimulation of thymidine uptake by 3T3 cells in the presence of heparin is ≤ 0.5 ng/ml corresponding to a specific activity of $\geq 2 \times 10^6$ units/mg.

Usage:

Angio-Proteomie's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Fax: (480) 247-4337, angioproteomie@gmail.com